



Water-Quality and Lake-Stage Data for Wisconsin Lakes, Water Year 2004



U.S. GEOLOGICAL SURVEY
Open-File Report 2005-1147

*Prepared in cooperation with the
State of Wisconsin and local agencies*



U.S. Department of the Interior
U.S. Geological Survey



Water-Quality and Lake-Stage Data for Wisconsin Lakes, Water Year 2004

A report by the Wisconsin District Lake-Studies Team—
W.J. Rose (team leader), H.S. Garn, G.L. Goddard, S.B. Marsh, D.L. Olson, and D.M. Robertson

Open-File Report 2005-1147



**Prepared in cooperation with the
State of Wisconsin and with other agencies**

Middleton, Wisconsin
2005

**U.S. DEPARTMENT OF THE INTERIOR
GALE A. NORTON, Secretary**

U.S. GEOLOGICAL SURVEY
Charles G. Groat, Director

For additional information write to:

Center Director
U.S. Geological Survey
8505 Research Way
Middleton, WI 53562

Copies of this report can be purchased from:

U.S. Geological Survey
Earth Science Information Center
Open-File Reports Section
Box 25286, MS 517
Denver Federal Center
Denver, CO 80225

CONTENTS

Introduction	1
Methods of data collection	4
Explanation of physical and chemical characteristics of lakes	9
Water temperature and thermal stratification	9
Specific conductance	11
Water clarity	12
pH	12
Dissolved oxygen	13
Phosphorus	14
Nitrogen	14
Chlorophyll <i>a</i>	15
Classification of lakes	15
References cited	17
Lake data	19
Big Cedar	
North Site, near West Bend, 432409088151600	21
South Site, near West Bend, 432224088154900	23
Butternut	
near Park Falls, 455854090310300	25
Deep Hole, near Park Falls, 455803090310800	26
North Site, near Butternut, 455904090303400	29
South Site, near Park Falls, 455731090311000	32
Far South Site, near Park Falls, 455651090312700	35
Delavan	
near Delavan, 423706088363400	37
at Center near Delavan Lake, 423556088365001	38
at North End near Lake Lawn, 423659088354401	44
at SW End near Delavan Lake, 423526088380101	44
Desair	45
Devils near Baraboo, 05404500	46
Fish near Sauk City, 05406050	47
Forest	48
Geneva	
at Lake Geneva, 423525088260400	50
West End, near Williams Bay, 423329088323300	51
Green	
at County Trunk Highway A near Green Lake (East End), 434928088553601	57
Deep Hole, 434756089020500	58
East End, 434928088570000	61
Koshkonong near Newville, 05427235	63

CONTENTS--Continued

Little Cedar	
North Site, near West Bend, 432255088134700	64
South Site, near West Bend, 432249088134500	66
Mendota at Madison, 05428000	68
Middle at Lauderdale, 424621088335500	69
Middle Genesee near Oconomowoc, 430309088284800	71
Monona at Madison, 05429000	74
Muskego (Big Muskego) near Wind Lake, 425109088075000	75
Nagawicka	
at Delafield, 430347088240800	76
at Deep Hole, at Delafield, 430417088230300	77
Oconomowoc	
No. 1 (Center) at Oconomowoc, 430551088273500	80
No. 2 (off Hewitt Point) at Oconomowoc, 430609088262200	82
Okauchee	
at Okauchee, 430723088252100	84
No. 1, near Okauchee, 430759088244200	86
No. 2, at Okauchee, 430645088264500	88
No. 3, at Okauchee, 430642088252400	90
No. 4, at Okauchee, 430757088261700	92
Potter near Mukwonago, 424905088204000	94
Powers at Powers Lake, 423246088175800	97
Spooner	
Deep Hole, near Spooner, 455034091493300	100
Southeast Site, near Spooner, 454945091483900	103
Tichigan near Waterford, 424854088123300	106
Twin	
East Twin, near Westfield, 435430089350700	109
West Twin, near Westfield, 435438089352300	111
Waterford Waterway, Fox River Site, near Waterford, 424614088123600	113
Wazee, Deep Hole, near Black River Falls, 441730090432900	116
Wind, at Outlet at Wind Lake, 424848088083100	118
at Wind Lake, 424915088083900	119
Winnebago	
at Oshkosh, 04082500	122
near Stockbridge, 04084255	123
Wisconsin District publications pertaining to lakes	124
Appendix - Quality-Assurance/Quality-Control Plan	133

CONTENTS--Continued

FIGURE

Figure 1. Map showing location of lake water-quality and lake-stage stations in Wisconsin 2

TABLES

Table 1. Discontinued lake stations	5
2. Parameter identification numbers and laboratory reporting levels (LRL) for chemical parameters commonly measured in lakes, and analyzed at the National Water-Quality Laboratory (NWQL) or the Wisconsin State Laboratory of Hygiene (WSLH)	11

CONVERSION FACTORS, VERTICAL DATUM, AND ABBREVIATED WATER-QUALITY UNITS

Multiply	By	To Obtain
mile (mi)	1.609	kilometer
pound (lb)	453.6	gram
acre	0.4048	hectare
foot (ft)	0.3048	meter
gallon (gal)	3.785	liter
square mile (mi^2)	2.590	square kilometer

Temperature, in degrees Celsius ($^{\circ}\text{C}$) can be converted to degrees Fahrenheit ($^{\circ}\text{F}$) by use of the following equation:

$$^{\circ}\text{F} = 1.8(^{\circ}\text{C}) + 32.$$

Sea level: In this report “sea level” refers to the National Geodetic Vertical Datum of 1929 (NGVD of 1929)—a geodetic datum derived from a general adjustment of the first-order level nets of both the United States and Canada, formerly called Sea Level Datum of 1929.

Abbreviated water-quality units: Chemical concentrations and water temperature are given in metric units. Chemical concentration is given in milligrams per liter (mg/L) or micrograms per liter ($\mu\text{g}/\text{L}$). Milligrams per liter is a unit expressing the concentration of chemical constituents in solution as weight (milligrams) of solute per unit volume (liter) of water. One thousand micrograms per liter is equivalent to one milligram per liter. For water with dissolved-solids concentrations less than 7,000 mg/L, the numerical values for concentrations expressed as mg/L and $\mu\text{g}/\text{L}$ are the same as for concentrations in parts per million and parts per billion, respectively.

Specific conductance of water is expressed in microsiemens per centimeter at 25 degrees Celsius ($\mu\text{S}/\text{cm}$). This unit is equivalent to micromhos per centimeter at 25 degrees Celsius ($\mu\text{mho}/\text{cm}$), formerly used by the U.S. Geological Survey.

WATER-QUALITY AND LAKE-STAGE DATA FOR WISCONSIN LAKES, WATER YEAR 2004

By Wisconsin Water Science Center Lake- Studies Team

INTRODUCTION

The U.S. Geological Survey (USGS), in cooperation with local and other agencies, collects data at selected lakes throughout Wisconsin. These data, accumulated over many years, provide a data base for developing an improved understanding of the water quality of lakes. To make these data available to interested parties outside the USGS, the data are published annually in this report series. The locations of water-quality and lake-stage stations in Wisconsin for water year 2004 are shown in figure 1. A water year is the 12-month period from October 1 through September 30. It is designated by the calendar year in which it ends. Thus, the period October 1, 2003 through September 30, 2004 is called "water year 2004."

The purpose of this report is to provide information about the chemical and physical characteristics of Wisconsin lakes. Data that have been collected at specific lakes, and information to aid in the interpretation of those data, are included in this report. Data collected include measurements of in-lake water quality and lake stage. Time series of Secchi depths, surface total phosphorus and chlorophyll *a* concentrations collected during non-frozen periods are included for all lakes. Graphs of vertical profiles of temperature, dissolved oxygen, pH, and specific conductance are included for sites where these parameters were measured. Descriptive information for each lake includes: location of the lake, area of the lake's watershed, period for which data are available, revisions to previously published records, and pertinent remarks. Additional data, such as streamflow and water quality in tributary and outlet streams of some of the lakes, are published in another volume: "Water Resources Data-Wisconsin, 2004."

Water-resources data, including stage and discharge data at most streamflow-gaging stations, are available through the World Wide Web on the Internet. The Wisconsin Water Science Center's home page is at <http://wi.water.usgs.gov/>. Information on the Wisconsin Water Science Center's Lakes Program is found at wi.water.usgs.gov/lake/index.html and wi.water.usgs.gov/projects/index.html.



Figure 1. Location of lake water-quality and lake-stage stations in Wisconsin.

The USGS has done cooperative lake monitoring with local and other agencies since 1983. Cooperators in 2004 included:

Big Cedar Lake Protection and Rehabilitation District
City of Delafield
City of Muskego
Dane County Department of Public Works
Delavan Lake Sanitary District
Geneva Lake Environmental Agency
Green Lake Sanitary District
Jackson County
Lauderdale Lakes Lake District
Little Cedar Lake Protection and Rehabilitation District
Middle Genesee Lake District
Okauchee Lake Management District
Potters Lake Protection and Rehabilitation District
Powers Lake District
Rock County Public Works Department
Spooner Lake District
Town of Auburn (Forest Lake Association)
Town of Rice Lake (Desair Lake Restoration, Inc.)
Town of Springfield (Twin Lakes Conservancy)
U.S. Army Corps of Engineers
Village of Oconomowoc Lake
Waterford Waterway Management District
Wind Lake Management District
Wisconsin Department of Natural Resources

Lake data-collection sites are identified by a unique identification number. Lake water-quality sites are identified by a 15-digit number that is a concatenation of the site's latitude, longitude, and a two-digit sequence number. The sequence number is used to distinguish between sites located at the same latitude-longitude designation. The site identification number is permanently assigned to the site; actual latitude and longitude of the site are subject to update and are stored separately. For some lakes, which have historical records of lake stage, an eight-to-ten digit number is assigned according to downstream order. Gaps are left in the numerical series to allow for new stations; hence, the numbers are not consecutive. The first two digits of the complete eight-to-ten digit number, such as 04087000 or 054310157, designate the major river basin. For example, "04" designates the St. Lawrence River Basin and "05" designates the Upper Mississippi River Basin.

The water-quality lake stations that were discontinued prior to water year 2004 are listed in table 1. Discontinued lake-stage stations are not included in this table.

This report is the culmination of a concerted effort by a number of people who collected, compiled, analyzed, verified, and organized the data, and who typed, edited, and assembled the report. The authors had primary responsibility for assuring that the information contained herein is accurate, complete, and adheres to USGS policy and established guidelines. Technicians in charge of the field offices are: T.J. Popowski (Rice Lake and Merrill), and S.A. March (Middleton). The data were collected and processed by S.M. Berg, C.J. Bloom, G.L. Goddard, D.E. Housner, S.B. Marsh, B.W. Olson, D.L. Olson, and J.G. Schuler. S.B. Marsh and P.A. Stark assembled, edited, and formatted the report. Additional assistance in preparation of the report was provided by M.M. Greenwood.

METHODS OF DATA COLLECTION

Depth profiles of water temperature, dissolved oxygen, pH, and specific conductance were collected using multi-parameter meters. Prior to measurements, the meters were calibrated using standards for pH and conductance, and dissolved oxygen was calibrated using the air calibration method. Generally, field measurements in profiles were made at 0.5-m intervals if the maximum depth of the lake was 5 m or less and at 1.0-m intervals if the maximum depth was greater than 5 m.

Table 1. Discontinued lake stations

Station name	Site identification number	Period of record
Alma Lake near St. Germain	455426089254700	Oct. 1984–Sept. 1990, May 1992–Sept. 1996
Balsam Lake, off Cedar Island, at Balsam Lake off Little Narrows, near Balsam Lake	452755092264600 452858092265300	Feb. 1991–Aug. 1994 May 1991–Aug. 1994
off Rock Island, near Balsam Lake	452754092234300	May 1991–Aug. 1994
Balsam Lake near Birchwood	453907091345800	Mar. 1993–Aug. 1994, Mar. 1996–Aug. 1997, Mar.–Sept. 2001
Bass Lake near Shawano	445215088300300	Feb. 1990–Aug. 1992
Bear Lake at Deep Hole near Haugen	453754091490900	Mar. 1992–Aug. 1993
Beaver Dam Lake, South end, at Beaver Dam North end, near Beaver Dam	432814088515000 433122088545700	June–Oct. 1991 June–Oct. 1991
Benedict Lake near Powers Lake	423201088180800	May 1998–Aug. 2000
Big Blacksmith Lake near Keshena	445401088334500	Feb. 1990–Aug. 1992
Big Hills (Hills) Lake near Wild Rose	440912089092000	June 1983–Aug. 1984, Feb.–Aug. 1987, Feb.–Aug. 1990, Feb.–Aug. 1993, Feb.–Aug. 1996, Feb.–Aug. 1999
Big Muskego Lake, at North Site, near Muskego Research Base, near Muskego	425301088061300 425235088075300	Feb.–Aug. 1988 May–June 1994
Big Round Lake near Milltown	453142092180100	Feb.–Sept. 2001
Big St. Germain Lake, near St. Germain near Lake Tomahawk	455557089311000 05390750	Feb. 1992–Aug. 1996 1991–2001
Big Sand Lake, Deep Hole, near Hertel East Site, near Hertel	454910092134000 454921092124300	Feb.–Sept. 2001 Feb.–Sept. 2001
Big Sissabagama Lake, near Stone Lake North Site, near Stone Lake	454724091303600 454800091312900	Apr. 1986–Sept. 1996 Oct. 1997–Sept. 2002 Mar. 1998–Sept. 2001
Booth Lake near East Troy	424800088254800	Feb. 1992–Aug. 1994 Feb. 2001–Aug. 2003
Buffalo Lake, Center Site, at Packwaukee East End, at Montello	434558089260600 434720089201600	May 1998–Sept. 2001 May 1998–Sept. 2001
West End, near Endeavor	434414089282400	May 1998–Sept. 2001
Denoon Lake at Wind Lake	425044088100300	Feb. 1991–Aug. 1996
Druid Lake near Hartford	431643088243300	Feb. 1991–Sept. 1996
Eagle Lake near Kansasville	05544500	1936–64, 1975–77, 1979, Feb. 1993–Sept. 1996
Eagle Lake, at Deep Hole, near Kansasville	424207088072400	Feb. 1993–Aug. 1996
Eagle Spring Lake at Eagleville	425103088261500	Apr. 1991–Sept. 2001
Elizabeth Lake near Twin Lakes	423051088155300	Feb. 1995–Sept. 1997
Fowler Lake, Center, at Oconomowoc	430653088294601	Jan.–Dec. 1984, Oct. 1986–Sept. 1996

Table 1. Discontinued lake stations

Station name	Site identification number	Period of record
Fox Lake Deep Hole at Fox Lake	433458088560600	June 1991–Mar. 1993
Geneva Lake		
Geneva Bay, at Lake Geneva	423455088263800	Apr. 1997–Feb. 1999
Williams Bay, at Williams Bay	423420088320500	Apr. 1997–Feb. 1999
Center, near Lake Geneva	423402088301400	Apr. 1997–Mar. 1999
East End, near Lake Geneva	423421088272300	Apr. 1997–May 2000
Hemlock Lake near Mikana	453421091333700	Mar. 1993–Aug. 1994, Mar. 1996–Aug. 1997, Mar.–Sept. 2001
Hooker Lake at Salem	423335088060300	Feb. 1992–Aug. 1993
Kawaguesaga		
Deep Hole, near Minocqua	455208089435800	May–Sept. 2003
South Site, near Minocqua	455145089442600	May–Sept. 2003
Kirby Lake near Cumberland		
(Site 1) near Cumberland	453554092042101	Nov. 1995–Oct. 1996
(Site 2) near Cumberland	453608092035801	Nov. 1995–Nov. 1996
(Site 3) near Cumberland	45361092035301	Nov. 1995–Nov. 1996
(Site 4) near Cumberland	453612092034901	Nov. 1995–Nov. 1996
(Site 5) near Cumberland	453603092035701	Nov. 1995–Nov. 1996
(Site 6) near Cumberland	453608092041201	Nov. 1995–Nov. 1996
Lac La Belle at Oconomowoc	430733088305900	Feb. 1984–Aug. 1985 Apr. –Aug. 1991 Feb. 2001–Aug. 2003
NW, at Oconomowoc	430809088313900	Feb. 1984–Aug. 1985
SE at Oconomowoc	430707088301400	Feb. 1984–Aug. 1985
Lake Blass at Lake Delton	433545089482400	Mar. 1989–Aug. 1990
Lake Keesus, East Bay, near Merton	430957088183400	Apr. 1991–Aug. 1995
North Bay, near Merton	431006088191000	Apr. 1991–Aug. 1995
Lake Morris at Mount Morris	440654089120500	Jun. 1983–Sept. 1989
Lake Nebagamon, Northeast Bay, at Lake Nebagamon	463050091412300	May 1992–Aug. 1995
Southeast Bay, at Lake Nebagamon	462928091413500	Mar. 1992–Sept. 1995
West Bay, at Lake Nebagamon	463034091425300	May 1992–Aug. 1995
Lake Noquebay near Crivitz	451511087550900	Feb. 1987–Aug. 1988, Apr. 1991–Aug. 1994
East End, near Crivitz	451540087525700	Apr. 1991–Aug. 1994
Lamotte Lake near Shawano	445305088361200	Feb. 1990–Aug. 1992
Lauderdale Lakes		
at Lauderdale	424554088332700	Oct. 1993–Oct. 1994
Green, Auxiliary, Number 1, near Lauderdale	424640088341900	June 1999–Sept. 2000
Green, near Lauderdale	424652088341500	Nov. 1993–Nov. 1994 Aug. 2002
Mill, at Lauderdale	424555088335700	Nov. 1993–Nov. 1994 Aug. 2002
Legend Lake (site 1) near Shawano	445342088312700	Feb. 1990–Feb. 1992

Table 1. Discontinued lake stations

Station name	Site identification number	Period of record
Little Arbor Vitae near Woodruff	455446089370300	Feb. 1991–Sept. 2002
Little Green Lake, at Center, near Markesan	434412088590700	Feb. 1991–Aug. 2003
Little Muskego Lake at Muskego	425425088083500	Oct. 1986–Aug. 2002
Little Rock Lake near Woodruff	455946089415702	Oct. 1983–Sept. 1996
Little St. Germain Lake near Eagle River	05390700	(a)
Upper East Bay, at St. Germain	455532089253900	Dec. 1996–Mar. 97
		Mar. 1999
		Mar. 2000–Aug. 2003
Northeast Bay, near St. Germain	455545089262500	Apr. 1991–Aug. 1994
		Aug. 1996–Aug. 1997
		Mar. 1999–Aug. 2003
South Bay, near St. Germain	455437089270800	Apr. 1991–Aug. 1994
		Aug. 1996–Aug. 1997
		Mar. 1999–Aug. 2003
West Bay, at St. Germain	455428089282400	Apr. 1991–Aug. 1994
		Aug. 1996–Aug. 1997
		Mar. 1999–Aug. 2003
Little Sand Lake - Site No. 2 - near Mole Lake	452826088544101	May 1996–Sept. 2003
Long (Kee Nong Go-Mong) Lake at Wind Lake	424937088103400	Feb. 1988–Aug. 1989, Feb. 1991–Aug. 1996
Loon Lake near Shawano	445009088303700	Feb. 1991–Aug. 1993
Lost Lake near Beaver Dam	432640088580500	June–Oct. 1991
McKenzie Lakes		
McKenzie (Big McKenzie)		
Deep Hole, near Spooner	455507092013500	Feb. 1987–Aug. 1998
Northern Site, near Spooner	455540092022000	June 1997–Aug. 1998
South Site, near Spooner	455437092022300	June 1997–Aug. 1998
Lower McKenzie, near Webb Lake	455902092011900	June 1997–Aug. 1998
Middle McKenzie, near Spooner	455635092021800	June 1997–Aug. 1998
Mary (Marie) Lake at Twin Lakes	423128088151200	Feb. 1995–Aug. 1997
Max Lake near Woodruff	460128089423501	Mar. 1988–Dec. 1996
Mead Lake, East Bay near Willard	444720090445000	Apr. 1991–Aug. 1995
West Bay near Willard	444733090460100	Feb. 1991–Sept. 1995
Minocqua Lake		
Deep Hole, at Minocqua	455214089412800	May–Sept. 2003
North Bay, at Minocqua	455232089424100	May–Sept. 2003
South Bay, at Minocqua	455206089425200	May–Sept. 2003
Montello Lake at Montello	434748089195800	Feb. 1995–Aug. 1998
Moon Lake near St. Germain	455504089260500	Feb. 1992–Aug. 1996
Morgan Lake near Fence	454622088324801	Oct. 1987–Sept. 1998.
Moshawquit Lake near Shawano	445352088295800	Feb. 1990–Aug. 1992
Muskego (Big Muskego)		
Auxiliary Number 1, near Muskego	425329088054000	June 1996–Aug. 2000

Table 1. Discontinued lake stations

Station name	Site identification number	Period of record
Bass Bay, near Muskego near Wind Lake	425344008807010 425109088075000	Feb. 1988–Aug. 2002 Oct. 1987–Sept. 1989, Jan. 1991–Sept. 2002
South Site, near Muskego	425212088072800	Feb. 1988–Aug. 2002
Muskellunge Lake near Eagle River	455700089224900	June 2000–Aug. 2001
Muskellunge Lake, near Lake Outlet near Eagle River near Eagle River	455706089232400 455700089224900	Nov. 2000–Oct. 2001 June 2000–Aug. 2001
Namekagon Lakes		
Garden, near Cable	461224091033200	Mar. 1998–Aug. 1999
Jackson, near Cable	461457091065900	Mar. 1998–Aug. 1999
Namekagon		
Deep Hole, near Cable	461308091065100	Mar. 1998–Aug. 1999
East Basin, near Cable	461228091044300	Mar. 1998–Aug. 1999
Northeast Basin, near Cable	461410091050700	Mar. 1998–Aug. 1999
Park Lake (site 1) at Pardeeville	433239089175800	Feb. 1986–Aug. 1987, May–Nov. 1993
(site 2) at Pardeeville	433226089175500	May–Nov. 1993
(site 3) at Pardeeville	433245089173000	May–Nov. 1993
(site 4) at Pardeeville	433257089165100	May–Nov. 1993
Pike Lake near Hartford	431916088200501	Dec. 1998–Dec. 2000
Pike Lake-QW Site-near Hartford	431835088200600	Feb.–Aug. 2000
Pretty Lake, at Deep Hole, near Dousman	425722088295000	Feb. 1993–Aug. 1997
Red Cedar Lake, at Mikana	453522091360600	Mar. 1993–Aug. 1994, Mar. 1996–Aug. 1997, Oct. 2000–Sept. 2001
Deep Hole, near Mikana	453725091345100	Mar. 1993–Aug. 1994, Mar. 1996–Aug. 1997, Mar. –Sept. 2001
South End, at Mikana	453519091352500	Mar. 1993–Aug. 1994, Mar. 1996–Aug. 1997, Mar. –Sept. 2001
Rice Lake at Deep Hole near Whitewater	424629088415700	Apr.–Nov. 1991
Round Lake near Shawano	445328088335000	Feb. 1990–Aug. 1992
Sand Lake (Deep Hole) near Keshena	445321088323101	June–Aug. 1992
Shell Lake at Shell Lake	05334000	Aug. 1936–Sept. 1999
Silver Lake near Oconomowoc	430436088293300	Apr. 1992–Aug. 1996
Silver Lake near West Bend	432322088125000	Feb. 1996–Aug. 1997
Sinissippi Lake, off Anthony Is., at Hustisford off Butternut Is., near Hustisford	432113088361100 432240088363900	Feb. 1991–Aug. 1993 Apr. 1991–Aug. 1993
off Sam Point, near Hustisford	432300088374200	Apr. 1991–Aug. 1993
Spirit Lake near Keshena	445400088320100	Apr.–Aug. 1992
Stewart Lake at Mt. Horeb	430117089442701	May 1992–Sept. 1993
Tombeau Lake near Powers Lake	423153088184800	May 1998–Aug. 2000
Upper Nemahbin Lake, Center, near Delafield	430400088254900	June 1993–Aug. 1995

Table 1. Discontinued lake stations

Station name	Site identification number	Period of record
South Site, near Delafield	430339088254800	June 1993–Aug. 1995
Outlet near Delafield	430334088255400	June 1993–Aug. 1995
Vandercook Lake near Woodruff	455909089405602	Nov. 1980–Aug. 1998
Watosah-skice Lake near Keshena	445330088361400	Feb. 1990–Aug. 1992
Waubeesee Lake at Wind Lake	424857088101500	Feb. 1988–Aug. 1989, Feb. 1991–Aug. 1996
Wazee Lake		
Beach, near Black River Falls	441721090431700	Nov. 1999–Aug. 2000 June–Sept. 2003
Whitefish Lake, North Basin, near Gordon	461321091520900	Mar. 1998–Sept. 2001
South Basin, near Gordon	461212091523200	Mar. 1998–Sept. 2001
Whitewater Lake, off Heart Prairie, near Whitewater	424533088420100	Apr.–Nov. 1991
near Whitewater	424608088414800	Nov. 1990–Sept. 2002
North Bay, near Whitewater	424625088405500	Apr.–Nov. 1991
South Bay, near Whitewater	424501088422300	Apr.–Nov. 1991
Wind Lake, Northeast Basin, at Wind Lake	424938088080800	Feb. 1997–Aug. 1998
Wolf Lake near Mt. Calvary	435152088123100	Nov. 1983–Sept. 1986, Nov. 1992–Sept. 1997

(a) Wisconsin Valley Improvement Co. currently collects stage data for this site.

In most lakes, water samples were collected at two depths - near the surface and near the bottom. Chemical analyses of water samples were performed using standard analytical methods by either the USGS National Water Quality Laboratory (Wershaw and others, 1987; Fishman and Friedman, 1989; Fishman, 1993) or the Wisconsin State Laboratory of Hygiene (Wisconsin State Laboratory of Hygiene, 1993). Analyses for dissolved constituents were performed on samples that were filtered in the field through a 0.45- μm (micrometer) pore-size filter. Total or total recoverable constituents were determined by analyzing unfiltered water samples. Preservation and shipment of samples followed standard protocols established by the laboratories. Water-quality data were archived in the Water Quality Data Base (QWDATA) of the National Water Information System (NWIS). Additional descriptive information about water-quality data is available in the data report: "Water Resources Data – Wisconsin, 2004". NWIS parameter codes and minimum laboratory reporting levels for chemical constituents are given in table 2.

Records of lake stage are considered complete when one or more manual or automatic measurements were obtained per day. Partial records of lake stage result when measurements were less frequent than daily. A complete description of manual or automatic measurements of lake stage is described by Rantz and others (1982).

EXPLANATION OF PHYSICAL AND CHEMICAL CHARACTERISTICS OF LAKES

Following are brief, generalized explanations of some of the common measurements of water quality and some of the physical processes occurring in lakes that influence these measures of water quality. More detailed explanations of water-quality data and lake processes are given by Wetzel (1983), Hem (1985), and Shaw and others (1993).

Water Temperature and Thermal Stratification

Water temperature in lakes is important because of its role in stratification and because of the temperature dependence of many chemical reactions and life processes of aquatic organisms. The extent of thermal stratification in lakes depends on the interaction between the lake's shape, water clarity, solar heating, and wind-driven mixing. Complete mixing of the lake is usually inhibited by thermal stratification in summer and by ice cover in winter. Thermal stratification affects water quality and the distribution of organisms in the lake. Summer thermal stratification can occur in any lake, but in Wisconsin it commonly occurs in lakes deeper than about 6 m (Shaw and others, 1993).

Table 2. Parameter identification numbers and laboratory reporting levels (LRL) for chemical parameters commonly measured in lakes, and analyzed at the National Water Quality Laboratory (NWQL) or the Wisconsin State Laboratory of Hygiene (WSLH)

Parameter Name	Units	CAS Number (1)	Parameter Code (2)	(NWQL)		(WSLH)		
				Standard Analysis		Low-Level Analysis		
				LRL	Lab Code	LRL	Lab Code	LRL
Calcium, diss. (Ca)	mg/L	7440-70-2	00915	0.020	659	0.002	1895	0.02
Magnesium, diss. (Mg)	mg/L	7439-95-4	00925	0.004	663	0.001	1897	0.02
Sodium, diss. (Na)	mg/L	7440-23-5	00930	0.09	675	0.025	1898	0.09
Potassium, diss. (K)	mg/L	7440-09-7	00935	0.24	54	0.01	833	0.3
Sulfate, diss. (SO ₄)	mg/L	14808-79-8	00945	0.31	1572	0.01	1263	1.0
Chloride, diss. (Cl)	mg/L	16887-00-6	00940	0.29	1571	0.01	1259	0.1
Fluoride, diss. (F)	mg/L	16984-48-8	00950	0.100	31	0.01	1260	0.03
Iron, diss. (Fe)	µg/L	7439-89-6	01046	10	645	3	1896	10
Manganese, diss. (Mn)	µg/L	7439-96-5	01056	2.2	648	1	1793	0.4
Silica, diss. (SiO ₂)	mg/L	7631-86-9	00955	0.1	56	0.02	1899	0.008
Nitrogen, NO ₂ +NO ₃ , diss.	mg/L		00631	0.05	1975	0.005	1979	0.01
Nitrogen, ammonia, diss.	mg/L	7664-41-7	00608	0.02	1976	0.002	1980	0.013
Nitrogen, organic, total (3)	mg/L							
Nitrogen, amm.+org., total (4)	mg/L	17778-88-0	00625	0.100	1985			0.2
Nitrogen, amm.+org.,diss.	mg/L		00623					I470DLD
Nitrogen, total (5)	mg/L							
Nitrogen, dissolved	mg/L		00602					
Phosphorus, total	mg/L	7723-14-0	00665	0.05	1984	0.004	2333	0.005
Phosphorus, ortho, diss.	mg/L	14265-44-2	00671	0.01	1262	0.002	1978	0.002
Chlorophyll a, phytoplankton	µg/L	479-61-8	70953	0.1	586			
Chlorophyll a, phytoplankton	µg/L	479-61-8	32210				0.26	I250UNF

Footnotes:

1: CAS (Chemical Abstracting Services) number = unique identification for each constituent

2: Parameter Code - unique number for storage of data in database

3: Calculated as difference between total ammonia + organic nitrogen and ammonia nitrogen

4: Also known as Total Kjeldahl Nitrogen (TKN)

5: Calculated as sum of TKN + Nitrogen as (NO₂+NO₃)

The density of water increases with decreasing temperature down to a temperature of 4°C, then decreases with decreasing temperature between 4°C and the freezing point of water (0°C). For a brief period in the spring after the ice is out, water temperature is usually uniform through the entire water column and wind action causes the lake to mix completely. This process is known as “spring turnover.” As the lake absorbs the sun’s energy, the surface water becomes warmer and its density decreases, making it more resistant to complete mixing. The difference in density caused by different water temperatures can prevent warm and cold water from mixing. In most lakes, therefore, a density “barrier” forms between the warmer surface water (epilimnion) and the underlying colder water (hypolimnion). This barrier is often marked by a sharp temperature gradient known as the “thermocline (metalimnion).” During the stratified summer period, these three distinct layers of lake water are often present. As the temperature difference between surface and deep water increases, this “stratified” condition stabilizes and can persist until surface temperatures decrease in the fall, which decreases the stability of the stratification. The mixing of the lake water in the fall is known as “fall turnover.”

Thermal stratification may also occur under ice cover in the winter. In the winter, the coldest water (near 0°C) under the ice at the surface of the lake is less dense than water deeper in the lake with warmer temperatures.

Specific Conductance

Specific conductance is a measure of the ability of water to conduct an electrical current and is an indicator of the concentration of dissolved solids in the water. Because conductance is temperature related, reported values are normalized at 25°C and are termed specific conductance. As the concentration of dissolved minerals increases, specific conductance increases. During winter and summer thermal stratification, concentrations of dissolved constituents near the lake bottom increase due to the decomposition of materials settling from the epilimnion, or release of dissolved materials (such as iron, manganese, and phosphorus) from the bottom sediments during anoxic periods. Therefore, differences in specific conductance with depth indicate differences in concentrations of dissolved solids.

Water Clarity

Water clarity, or transparency, is commonly measured using a Secchi disc. The range of depths within which photosynthetic activity occurs depends largely on depth of light penetration, which is influenced by water clarity. A Secchi disc, most commonly an 20-cm.-diameter disc with alternating black-and-white quadrants, is lowered to a depth at which it is no longer visible. This depth is referred to as the Secchi depth. Clarity can be reduced by algae, zooplankton, water color, and suspended sediment. Algae are often the most dominant influence on clarity in lakes and, therefore, Secchi depth is usually correlated with the algal abundance. Secchi depths are generally the least during summer when algal populations are largest.

pH

The pH is a measure of the acidity of the water. It is defined as the negative logarithm of hydrogen-ion concentration and varies over a 14-unit log scale, with a pH of 7 being neutral. Values less than 7 indicate acidic conditions; the lower the value, the stronger the acidity. Values greater than 7 indicate alkaline conditions. The pH of water is influenced in part by photosynthesis and respiration of planktonic algae and aquatic plants. It is important because it affects the solubility of many chemical constituents, and because aquatic organisms have limited pH tolerances. Planktonic algae and aquatic plants produce oxygen and consume carbon dioxide as they photosynthesize during daytime; they consume oxygen and produce carbon dioxide when they respire at night. Carbon dioxide combines with the water molecule to form carbonic acid; therefore respiration causes a decrease in pH at night and photosynthesis during the day causes an increase in pH. The result is a daily cycle in pH. Because phytoplankton are usually concentrated in the near-surface water, changes in pH in the epilimnion are more extreme than in the hypolimnion, where less photosynthesis usually occurs.

Lakes having good fish populations and productivity generally have a pH between 6.7 and 8.2. Values of pH greater than 8.5 have been shown to cause the release of phosphorus from lake sediments (James and Barko, 1991).

Dissolved Oxygen

Dissolved oxygen is one of the most critical factors affecting a lake ecosystem because it is essential to most aquatic organisms, and it is involved in many chemical reactions. Very low dissolved oxygen concentrations can control some types of chemical reactions. The solubility of oxygen in water is inversely related to temperature—that is, oxygen solubility decreases as water temperature increases. This relation is important because at warmer temperatures the metabolic rate of organisms increases but less oxygen is available for respiration. The primary sources of dissolved oxygen are from the air and from photosynthesis. The minimum dissolved oxygen concentration specified in national water-quality criteria for early life stages of warmwater aquatic life is 5.0 mg/L (U.S. Environmental Protection Agency, 1986).

In early summer, if thermal stratification develops, the metalimnion restricts the surface supply of dissolved oxygen to the hypolimnion. The hypolimnion can become isolated from the atmosphere. Thus, as summer progresses, the dissolved oxygen concentration can decrease in response to decomposition of dead algae that settle from the epilimnion and in response to the biological and chemical oxygen demand of the sediments. The oxygen demand from these processes may completely deplete the oxygen (anoxia) in the water near the lake bottom. The oxygen depletion then progresses upward but usually is confined to the hypolimnion.

Anoxia in the hypolimnion is common in stratified eutrophic (nutrient-rich) lakes in Wisconsin. Complete anoxia, however, is often not detected because of meter constraints. During anoxic conditions, many aquatic organisms cannot survive, but many other species (primarily bacteria) actually function only in such conditions. Therefore, a shift from oxic to anoxic conditions produces a rapid and dramatic change in the biological community and chemical environment. Anoxia also can cause release of phosphorus from the bottom sediments. This phosphorus then mixes throughout the water column during spring and fall turnover.

Phosphorus

Phosphorus is one of the essential nutrients for plant growth. High phosphorus concentrations can cause dense algal populations (blooms) and can therefore be a major cause of eutrophication in lakes. When phosphorus concentrations exceed 0.025 mg/L at the time of spring overturn in lakes and reservoirs, these water bodies may occasionally experience excess or nuisance growth of algae or other aquatic plants (U.S. Environmental Protection Agency, 1986). In many regions of the country, including the upper Midwest, other nutrients, particularly nitrogen, tend to be in abundant supply. Phosphorus is often the nutrient in shortest supply, therefore limiting or controlling plant growth. About 90 percent of the lakes in Wisconsin are limited by phosphorus (Shaw and others, 1993). In water, dissolved orthophosphate is that part of total phosphorus that is most readily available for use by algae.

Internal phosphorus recycling occurs in many lakes. Phosphorus used by algae, aquatic plants, fish, and zooplankton is stored within these organisms. As these organisms die and decompose, this phosphorus is returned to the lake water and sediments. Anoxia in the hypolimnion makes phosphorus more soluble, adding further to the release of phosphorus from the falling particles and the lake sediments. During spring and fall turnover the phosphorus, which was released from the bottom sediments into the hypolimnion during anoxia, is mixed throughout the lake. The phosphorus is then available for algal growth. These phenomena are part of the internal-recycling processes of lakes.

Nitrogen

Nitrogen, like phosphorus, is an essential nutrient for plant and algal growth. Usually in Wisconsin lakes, nitrogen is in abundant supply from the atmosphere and other sources. If phosphorus is abundant relative to algal needs, nitrogen can become the limiting nutrient. In that case, algal blooms are more likely to be triggered by increases in nitrogen than by increases in phosphorus. Some bluegreen algal species can fix nitrogen from the atmosphere (Wetzel, 1983). Therefore, in situations where other types of algae are excluded because of a shortage of nitrogen, the nitrogen-fixing bluegreen algae have a competitive advantage and may be present in abundance.

Lakes with a nitrogen to phosphorus ratio larger than 15 to 1 near the surface may generally be considered phosphorus limited; a ratio from 10 to 1 to 15 to 1 indicates a transition situation; and a ratio smaller than 10 to 1 generally indicates nitrogen limitation. Total nitrogen is the sum of ammonia, organic nitrogen, and nitrate-plus-nitrite nitrogen. The near-surface concentration is commonly used to compute the total nitrogen to phosphorus ratio because most algal species grow near the lake surface.

Chlorophyll a

Chlorophyll *a* is a photosynthetic pigment found in algae (Wetzel, 1983) and other green plants. Its concentration, therefore, is commonly used as a measure of the density of the algal population in a lake. Chlorophyll *a* concentrations are generally highest during summer when algal populations are highest. Moderate populations of desirable algae are important in the food chain; however, excessive populations or algal blooms are undesirable. Algal blooms can cause taste and odor problems, and limit light penetration needed to support growth of submerged aquatic plants. Certain species of bluegreen algae can produce toxins (Rapavich and others, 1987).

CLASSIFICATION OF LAKES

Two methods are commonly used to classify and evaluate Wisconsin lakes according to their water quality or trophic state: Lillie and Mason's (1983) water-quality index and a modification of Carlson's (1977) Trophic State Index (TSI) by Lillie and others (1993). Three water-quality measures are used in these classification systems: near-surface concentrations of total phosphorus and chlorophyll *a*, and water clarity as indicated by the Secchi depth.

Lillie and Mason's (1983) water-quality indices for Wisconsin lakes were developed based on random summer measurements of total phosphorus and chlorophyll *a* concentrations, and Secchi depth to classify the lakes' water quality as shown below:

Water-quality index	Total phosphorus range (mg/L)	Chlorophyll <i>a</i> range ($\mu\text{g/L}$)	Water clarity range (Secchi depth, in meters)
"Excellent"	<0.001	<1.0	>6.0
"Very good"	.001–.009	1.0– 4.9	3.0–6.0
"Good"	.010–.029	5.0– 9.9	2.0–2.9
"Fair"	.030–.049	10.0–14.9	1.5–1.9
"Poor"	.050–.149	15.0–30.0	1.0–1.4
"Very poor"	>.150	>30.0	<1.0

The TSI approach to lake classification assigns numerical ranges to the three trophic conditions generally used to describe the wide range of lake water-quality conditions. Oligotrophic lakes are typically clear, algal populations and phosphorus concentrations are low, and the deepest water is likely to contain oxygen throughout the year. Mesotrophic lakes typically have a moderate supply of nutrients, experience moderate algal blooms, and have occasional oxygen depletions at depth. Eutrophic lakes are nutrient rich with relatively severe water-quality problems, such as frequent seasonal algal blooms, oxygen depletion in lower parts of the lakes, and poor clarity. When eutrophic conditions are very severe, the lake is considered hypereutrophic.

The WDNR modified the lakes classification scheme developed by Carlson (1977) to apply specifically to Wisconsin lakes. The WDNR system (Lillie and others, 1993) uses surface total phosphorus and chlorophyll *a* concentrations, and Secchi depth for ice-free periods to calculate values for TSI's. The WDNR has adopted the following TSI ranges to classify Wisconsin lakes: indices of less than 40 define oligotrophic conditions, 40 to 50 define mesotrophic conditions, greater than 50 to define eutrophic conditions, and greater than 70 define hypereutrophic conditions (Wisconsin Department of Natural Resources, 1992). These ranges are used to make relative comparisons in Wisconsin lake trophic-state evaluations by the WDNR and others.

The TSI for a lake can be calculated using the following equations (Lillie and others, 1993):

$$TSI_{\text{Secchi}} = 60.0 - 33.2 \times (\log_{10} \text{Secchi depth})$$

$$TSI_{\text{chlorophyll } a} = 34.82 + (17.41 \times (\log_{10} \text{chlorophyll } a \text{ concentration}))$$

$$TSI_{\text{total phosphorus}} = 28.24 + (17.81 \times (\log_{10} [\text{total phosphorus concentration} \times 1,000])))$$

where: Secchi depth is in meters,
chlorophyll *a* is in micrograms per liter, and
total phosphorus is in milligrams per liter.

The three trophic conditions are defined with the following boundaries for total phosphorus, Secchi disc, and chlorophyll *a*:

Trophic Level	Trophic State Index	Total phosphorus (mg/L)	Secchi depth (m)	Chlorophyll <i>a</i> ($\mu\text{g}/\text{L}$)
Eutrophic	50	0.017	2.0	7.4
Mesotrophic	40	0.005	4.0	2.0
Oligotrophic				

REFERENCES CITED

- Carlson, R.E., 1977, A trophic state index for lakes: Limnology and Oceanography, March, v. 22, no. 2, p. 361-369.
- Fishman, M.J., ed., 1993, Methods of analysis by the U.S. Geological Survey National Water Quality Laboratory—Determination of inorganic and organic constituents in water and fluvial sediments: U.S. Geological Survey Open-File Report 93-125, 217 p.
- Fishman, M.J., and Friedman, L.C., eds., 1989, Methods for determination of inorganic substances in water and fluvial sediments (3d ed.): U.S. Geological Survey Techniques of Water-Resources Investigations, book 5, chap. A1, 545 p.
- Hem, J.D., 1985, Study and interpretation of the chemical characteristics of natural water (3rd ed.): U.S. Geological Survey Water-Supply Paper 2254, 263 p.
- James, W.F., and Barko, J.W., 1991, Littoral-pelagic phosphorus dynamics during nighttime convective circulation: Limnology and Oceanography, v. 36, no. 5, p. 946-960.
- Lillie, R.A., Graham, S., and Rasmussen, P., 1993, Trophic-State Index equations and regional predictive equations for Wisconsin lakes: Wisconsin Department of Natural Resources Research Management Findings No. 35, 4 p.
- Lillie, R.A., and Mason, J.W., 1983, Limnological characteristics of Wisconsin lakes: Wisconsin Department of Natural Resources Technical Bulletin No. 138, 116 p.
- Rantz, S.E., and others, 1982, Measurement and computation of streamflow: U.S. Geological Survey Water-Supply Paper 2175, 631 p.
- Rapavich, W.M., Sonzogni, W.C., Standridge, J.H., Vennie J.G., and Wedepohl, R.E., 1987, Incidence of algal toxins in Wisconsin water experiencing blue-green algae blooms: Wisconsin State Laboratory of Hygiene and Wisconsin Department of Natural Resources, Informational Paper, 8 p.
- Shaw, B., Mechenich, C., and Klessig, L., 1993, Understanding Lake Data: G3582: University of Wisconsin Extension, Madison, Wis., 19 p.
- U.S. Environmental Protection Agency, 1986, Quality Criteria for Water 1986: U.S. Environmental Protection Agency publication, EPA 440/5-86-001 [variously paged].
- Wershaw, R.L., Fishman, M.J., Grabbe, R.R., and Lowe, L.E., eds., 1987, Methods for the determination of organic substances in water and fluvial sediments: U.S. Geological Survey Techniques of Water-Resources Investigations, book 5, chap. A3, 80 p.
- Wetzel, R.G., 1983, Limnology (2d ed.): New York, W.B. Saunders, 767 p.

Wisconsin Department of Natural Resources, 1992, Wisconsin water quality assessment—Report to Congress, 1992: Wisconsin Department of Natural Resources Publ-WR254-92-REV, 220 p.

Wisconsin State Laboratory of Hygiene, Environmental Sciences Section, 1993, Manual of analytical methods, inorganic chemistry unit: Wisconsin State Laboratory of Hygiene, revised November 1993 [variously paged].

LAKE DATA

432409088151600 BIG CEDAR LAKE, NORTH SITE, NEAR WEST BEND, WI

LOCATION.--Lat 43°24'09", long 88°15'16", in NE 1/4 sw 1/4 sec.20, T.11 N., R.19 E., Washington County, Hydrologic Unit 04040003, near West Bend.


PERIOD OF RECORD.--February 2000 to to current year.


REMARKS.--Lake sampled on north side at a depth of 12 m. Lake ice-covered during February sampling. Water-quality analyses by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, FEBRUARY 24 TO AUGUST 18, 2004

(Milligrams per liter unless otherwise indicated)

Date	<u>Feb-24</u>	<u>Apr-22</u>	<u>Jun-9</u>	<u>Jul-14</u>	<u>Aug-18</u>
Secchi depth (m)	--	1.4	3.1	1.1	1.45
Depth of sample (m)	0.5	11	0.5	11.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	--	15.2	--	9.5
Water temperature ($^{\circ}\text{C}$)	0.8	3.2	12.1	11.5	24.1
Specific conductance ($\mu\text{S/cm}$)	597	749	550	549	516
pH	7.8	7.1	8.3	8.2	7.5
Dissolved oxygen (mg/L)	12.6	3.5	10.6	9.9	10.6
Phosphorus, total (as P)	0.013	0.018	0.032	0.042	0.015
				0.026	0.014
				0.034	0.013
					0.040





Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Big Cedar Lake, North Site, near West Bend, Wisconsin.

432224088154900 BIG CEDAR LAKE, SOUTH SITE, NEAR WEST BEND, WI


LOCATION.--Lat 43°22'24", long 88°15'49", in NE 1/4 SE 1/4 sec.31, T.11 N., R.19 E., Washington County, Hydrologic Unit 04040003, near West Bend.


PERIOD OF RECORD.--February 2000 to current year.

REMARKS.--Lake sampled on south side at deep hole. Lake ice-covered during February sampling. Water-quality analyses by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, FEBRUARY 24 TO AUGUST 18, 2004
(Milligrams per liter unless otherwise indicated)

Date	<u>Feb-24</u>	<u>Apr-22</u>	<u>Jun-9</u>	<u>Jul-14</u>	<u>Aug-18</u>
	--	3.3	7.9	1.6	1.8
Secchi depth (m)					
Depth of sample (m)	0.5	30	0.5	30	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	--	8.8	--	1.49
Water temperature ($^{\circ}\text{C}$)	0.9	2.6	8.3	6.1	21.8
Specific conductance ($\mu\text{S/cm}$)	570	596	549	550	519
pH	7.9	7.4	8.3	8.1	8.3
Dissolved oxygen (mg/L)	13.5	1.1	12.1	11.4	11.1
Phosphorus, total (as P)	0.017	0.083	0.017	0.019	0.010
Phosphorus, ortho, dissolved (as P)	--	--	0.002	--	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	--	--	0.124	--	<0.019
Nitrogen, ammonia, dissolved (as N)	--	--	0.072	--	<0.015
Nitrogen, ammon. + diss., total (as N)	--	--	--	--	0.62
Nitrogen, amm. + org., total (as N)	--	--	0.49	--	--
Nitrogen, total (as N)	--	--	0.61	--	--
Color (Pt-Co. scale)	--	--	5	--	--
Turbidity (NTU)	--	--	1.9	--	--
Hardness, as CaCO_3	--	--	230	--	--
Calcium, dissolved (Ca)	--	--	36	--	--
Magnesium, dissolved (Mg)	--	--	33.7	--	--
Sodium, dissolved (Na)	--	--	20.4	--	--
Potassium, dissolved (K)	--	--	2	--	--
Alkalinity, as CaCO_3	--	--	192	--	--
Sulfate, dissolved (SO_4)	--	--	20.5	--	--
Chloride, dissolved (Cl)	--	--	44.3	--	--
Silica, dissolved (SiO_2)	--	--	0.389	--	--
Solids, dissolved, at 180°C	--	--	306	--	--
Iron, dissolved (Fe) ($\mu\text{g/L}$)	--	--	<100	--	--
Manganese, dissolved, (Mn) ($\mu\text{g/L}$)	--	--	<1	--	--





Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Big Cedar Lake, South Site, near West Bend, Wisconsin.

455854090310300 BUTTERNUT LAKE NEAR PARK FALLS, WI

LOCATION.--Lat 45°58'54", long 90°31'03", in NW 1/4 NW 1/4 NE 1/4 sec.5, T.40 N., R.1 W., Ashland County, Hydrologic Unit 07050002, about 150 ft south of Wisconsin Department of Natural Resources boat landing off County Highway B about 5 mi northwest of Park Falls.

DRAINAGE AREA.--47.6 mi², at lake outlet. Area of Butternut Lake is 1.57 mi².

PERIOD OF RECORD.--October 2002 to October 2004.

GAGE.--Water-stage recorder. Datum of gage is about 1,495 ft above sea level.

EXTREMES FOR PERIOD OF RECORD (NOVEMBER 2002 TO OCTOBER 2004).--Maximum recorded gage height, 7.38 ft, May 13, 2003; minimum recorded, 4.79 ft, July 30, 2003.

EXTREMES FOR CURRENT PERIOD (OCTOBER 2003 TO OCTOBER 2004).--Maximum recorded gage height, 7.17 ft, April 21; minimum recorded, 4.88 ft, July 29.

GAGE HEIGHT, FEET
WATER YEAR OCTOBER 2003 TO OCTOBER 2004
DAILY MEAN VALUES

DAY	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT
1	5.12	5.27	5.12	5.03	4.95	4.99	6.06	5.49	5.71	5.01	5.01	5.11	5.01
2	5.12	5.27	5.10	5.03	4.99	5.01	6.06	5.42	5.76	5.00	5.01	5.12	5.06
3	5.12	5.26	5.08	5.04	5.01	5.02	6.07	5.37	5.72	4.99	5.00	5.12	5.07
4	5.12	5.27	5.07	5.03	5.00	5.01	6.04	5.31	5.65	5.04	4.96	5.13	5.07
5	5.12	5.26	5.06	5.03	4.99	5.04	6.00	5.28	5.60	5.06	4.95	5.24	5.08
6	5.12	5.25	5.05	5.02	5.00	5.05	5.99	5.22	5.57	5.06	4.93	5.43	5.09
7	5.11	5.23	5.04	5.01	5.00	5.07	5.99	5.19	5.54	5.08	4.93	5.44	5.10
8	5.12	5.21	5.04	5.00	4.99	5.06	6.03	5.16	5.51	5.07	4.96	5.44	5.13
9	5.13	5.19	5.06	4.99	4.99	5.05	6.04	5.16	5.49	5.06	4.99	5.40	5.13
10	5.13	5.17	5.08	4.98	4.99	5.05	6.04	5.18	5.45	5.05	5.03	5.36	5.13
11	5.16	5.16	5.07	4.97	4.99	5.07	5.99	5.17	5.40	5.05	5.06	5.31	5.13
12	5.21	5.17	5.06	4.97	4.99	5.06	5.90	5.18	5.39	5.06	5.08	5.26	5.14
13	5.21	5.16	5.05	4.97	4.99	5.07	5.80	5.23	5.37	5.06	5.10	5.22	5.14
14	5.22	5.15	5.04	4.99	4.99	5.08	5.70	5.38	5.33	5.05	5.10	5.17	5.15
15	5.22	5.15	5.03	4.99	4.97	5.06	5.62	5.47	5.30	5.04	5.10	5.24	5.15
16	5.21	5.16	5.04	4.99	4.96	5.05	5.57	5.50	5.26	5.01	5.11	5.26	5.16
17	5.20	5.19	5.03	5.00	4.95	5.05	5.54	5.48	5.23	4.98	5.12	5.26	5.16
18	5.20	5.29	5.03	4.99	4.95	5.04	5.64	5.44	5.20	4.97	5.12	5.25	5.16
19	5.20	5.34	5.02	4.98	4.95	5.03	6.25	5.40	5.17	4.96	5.09	5.23	5.16
20	5.20	5.37	5.02	4.98	4.99	5.03	6.96	5.35	5.14	5.00	5.07	5.20	5.17
21	5.19	5.36	5.01	4.99	4.99	5.01	7.11	5.31	5.11	5.03	5.06	5.16	5.18
22	5.18	5.33	5.00	4.98	4.98	5.00	6.89	5.28	5.09	5.00	5.05	5.13	5.20
23	5.16	5.32	5.00	4.98	4.99	4.99	6.60	5.29	5.09	4.97	5.05	5.11	5.28
24	5.17	5.29	5.00	4.98	4.99	4.99	6.32	5.37	5.09	4.97	5.05	5.10	5.34
25	5.18	5.27	4.99	4.98	4.98	5.00	6.13	5.44	5.09	4.95	5.06	5.08	5.37
26	5.18	5.23	4.99	4.97	4.98	5.04	5.98	5.47	5.07	4.93	5.07	5.06	5.38
27	5.18	5.20	4.99	4.98	4.97	5.11	5.86	5.49	5.07	4.92	5.08	5.04	5.39
28	5.21	5.18	5.01	4.97	4.97	5.31	5.76	5.47	5.05	4.91	5.08	5.02	5.42
29	5.22	5.16	5.02	4.96	4.97	5.65	5.66	5.45	5.04	4.94	5.08	5.00	5.50
30	5.25	5.14	5.03	4.96	---	5.89	5.58	5.46	5.03	4.95	5.10	4.99	5.58
31	5.27	---	5.04	4.95	---	6.01	---	5.59	---	5.00	5.10	---	5.61
MEAN	5.18	5.23	5.04	4.99	4.98	5.13	6.04	5.35	5.32	5.01	5.05	5.20	5.21
MAX	5.27	5.37	5.12	5.04	5.01	6.01	7.11	5.59	5.76	5.08	5.12	5.44	5.61
MIN	5.11	5.14	4.99	4.95	4.95	4.99	5.54	5.16	5.03	4.91	4.93	4.99	5.01

455803090310800 BUTTERNUT LAKE, DEEP HOLE, NEAR PARK FALLS, WI

LOCATION.--Lat 45°58'03", long 90°31'08", in NW 1/4 NW 1/4 NE 1/4 sec.8, T.40 N., R.1 W., Ashland County, Hydrologic Unit 07050002, near Park Falls.


PERIOD OF RECORD.--March 2003 to current year.

REMARKS.--Lake sampled at deep hole at a lake depth of about 8 m. Lake ice-covered during March sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, MARCH 17 TO JUNE 2, 2004
(Milligrams per liter unless otherwise indicated)


Date	Mar-17	May-4	Jun-2
Lake stage (ft)	5.05	5.31	5.76
Secchi depth (m)	--	1.0	1.3
Depth of sample (m)	1	7	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	7.4	3.4
Water temperature ($^{\circ}\text{C}$)	0	4.8	14.7
Specific conductance ($\mu\text{S/cm}$)	133	207	79
pH	7.0	7.0	7.2
Dissolved oxygen (mg/L)	8.4	0.5	8.0
Phosphorus, total (as P)	0.030	0.309	0.038
Phosphorus, ortho, dissolved (as P)	--	0.011	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	--	0.133	--
Nitrogen, ammonia, dissolved (as N)	--	0.019	--
Nitrogen, amm. + org., total (as N)	--	0.75	--
Nitrogen, total (as N)	--	0.88	--
Color (Pt-Co. scale)	--	100	--
Turbidity (NTU)	--	8.9	--
Hardness, as CaCO_3	--	37	--
Calcium, dissolved (Ca)	--	9.6	--
Magnesium, dissolved (Mg)	--	3.1	--
Sodium, dissolved (Na)	--	2.1	--
Potassium, dissolved (K)	--	1	--
Alkalinity, as CaCO_3	--	30	--
Sulfate, dissolved (SO_4)	--	5.4	--
Chloride, dissolved (Cl)	--	3.2	--
Silica, dissolved (SiO_2)	--	8.88	--
Solids, dissolved, at 180°C	--	70	--
Iron, dissolved (Fe) ($\mu\text{g/L}$)	--	300	--
Manganese, dissolved, (Mn) ($\mu\text{g/L}$)	--	40	--

3-17-04




DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER


6-2-04



WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS



PH IN STANDARD UNITS





SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEGMENS PER CENTIMETER AT 25 DEGREES CELSIUS

455803090310800 BUTTERNUT LAKE, DEEP HOLE, NEAR PARK FALLS, WI--CONTINUED

WATER-QUALITY DATA, JULY 21 TO SEPTEMBER 21, 2004
 (Milligrams per liter unless otherwise indicated)

Date	<u>Jul-21</u>	<u>Aug-10</u>	<u>Sep-21</u>
Lake stage (ft)	5.03	5.03	5.16
Secchi depth (m)	0.9	1.0	1.5
Depth of sample (m)	0.5	9	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	35.7	--	--
Water temperature ($^{\circ}\text{C}$)	25.0	18.3	18.7
Specific conductance ($\mu\text{S/cm}$)	91	100	102
pH	8.6	6.5	7.7
Dissolved oxygen (mg/L)	10.9	0.6	8.4
Phosphorus, total (as P)	0.044	0.059	0.057
Phosphorus, ortho, dissolved (as P)	0.002	--	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	0.019	--	--
Nitrogen, ammonia, dissolved (as N)	0.025	--	--
Nitrogen, amm. + diss., total (as N)	0.82	--	--





Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Butternut Lake, Deep Hole, near Park Falls, Wisconsin.

455904090303400 BUTTERNUT LAKE, NORTH SITE, NEAR BUTTERNUT, WI


LOCATION.--Lat 45°59'04", long 90°30'34", in NE 1/4 SE 1/4 NE 1/4 sec.32, T.41 N., R.1 W., Ashland County, Hydrologic Unit 07050002, near Butternut.

PERIOD OF RECORD.--March 2004 to current year.

REMARKS.--Lake sampled at North Site at a lake depth of about 5.5 m. Lake ice-covered during March sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, MARCH 17 TO JUNE 2, 2004
(Milligrams per liter unless otherwise indicated)


Date	Mar-17	May-4	Jun-2
Lake stage (ft)	5.05	5.31	5.76
Secchi depth (m)	--	1.1	1.3
Depth of sample (m)	1	5	5.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	4.1	--
Water temperature ($^{\circ}\text{C}$)	-0.1	3.0	9.6
Specific conductance ($\mu\text{S/cm}$)	140	163	66
pH	7.0	6.8	6.4
Dissolved oxygen (mg/L)	8.0	4.5	8.7
Phosphorus, total (as P)	0.030	0.034	0.046
			0.040
			0.044
			0.040




455904090303400 BUTTERNUT LAKE, NORTH SITE, NEAR BUTTERNUT, WI--CONTINUED

WATER-QUALITY DATA, JULY 21 TO SEPTEMBER 21, 2004
 (Milligrams per liter unless otherwise indicated)

Date	<u>Jul-21</u>	<u>Aug-10</u>	<u>Sep-21</u>
Lake stage (ft)	5.03	5.03	5.16
Secchi depth (m)	1.0	1.0	1.1
Depth of sample (m)	0.5	5.8	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	34.6	--	--
Water temperature ($^{\circ}\text{C}$)	25.2	17.9	18.9
Specific conductance ($\mu\text{S/cm}$)	94	110	104
pH	8.7	6.6	7.7
Dissolved oxygen (mg/L)	11.3	0.2	7.6
Phosphorus, total (as P)	0.045	0.042	0.057
			0.067
			0.063





Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Butternut Lake, North Site, near Butternut, Wisconsin.

455731090311000 BUTTERNUT LAKE, SOUTH SITE, NEAR PARK FALLS, WI


LOCATION.--Lat 45°57'31", long 90°31'10", in SW 1/4 NW 1/4 SE 1/4 sec.8, T.40 N., R.1 W., Ashland County, Hydrologic Unit 07050002, near Park Falls.

PERIOD OF RECORD.--March 2003 to current year.

REMARKS.--Lake sampled at South Site at a lake depth of about 6 m. Lake ice-covered during March sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, MARCH 17 TO JUNE 2, 2004
(Milligrams per liter unless otherwise indicated)


Date	Mar-17	May-4	Jun-2
Lake stage (ft)	5.05	5.31	5.76
Secchi depth (m)	--	1.2	1.2
Depth of sample (m)	1	6	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	6.3	5.6
Water temperature ($^{\circ}\text{C}$)	0.3	4.6	14.2
Specific conductance ($\mu\text{S/cm}$)	127	80	79
pH	6.8	6.7	7.2
Dissolved oxygen (mg/L)	7.5	0.8	8.0
Phosphorus, total (as P)	0.038	0.088	0.054
		0.047	0.034
		0.048	




455731090311000 BUTTERNUT LAKE, SOUTH SITE, NEAR PARK FALLS, WI--CONTINUED

WATER-QUALITY DATA, JULY 21 TO SEPTEMBER 21, 2004
 (Milligrams per liter unless otherwise indicated)

Date	<u>Jul-21</u>	<u>Aug-10</u>	<u>Sep-21</u>
Lake stage (ft)	5.03	5.03	5.16
Secchi depth (m)	1.1	1.0	1.3
Depth of sample (m)	0.5	0.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	33.2	--	20.8
Water temperature ($^{\circ}\text{C}$)	24.3	17.5	18.6
Specific conductance ($\mu\text{S/cm}$)	91	111	102
pH	8.4	6.6	7.5
Dissolved oxygen (mg/L)	10.3	0.2	8.3
Phosphorus, total (as P)	0.045	0.101	0.055
		0.047	0.051
		0.050	





Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Butternut Lake, South Site, near Park Falls, Wisconsin.

455651090312700 BUTTERNUT LAKE, FAR SOUTH SITE, NEAR PARK FALLS, WI


LOCATION.--Lat 45°56'51", long 90°31'27", in SW 1/4 SE 1/4 NE 1/4 sec.17, T.40 N., R.1 W., Ashland County, Hydrologic Unit 07050002, near Park Falls.


PERIOD OF RECORD.--March 2003 to current year.

REMARKS.--Lake sampled at deep hole at Far South Site at a lake depth of about 4 m. Lake ice-covered during March sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, JUNE 2 TO SEPTEMBER 21, 2004
(Milligrams per liter unless otherwise indicated)

Date	Jun-2	Jul-21	Aug-10	Sep-21
Lake stage (ft)	5.76	5.03	5.03	5.16
Secchi depth (m)	1.1	1.0	1.2	1.5
Depth of sample (m)	0.5	4.2	4	0.5
Water temperature (°C)	14.7	14.3	23.7	18.5
Specific conductance ($\mu\text{S}/\text{cm}$)	79	79	90	101
pH	7.1	7.3	8.2	7.0
Dissolved oxygen (mg/L)	8.4	8.3	10.2	7.9





Secchi depths, and TSI data for Butternut Lake, Far South Site, near Park Falls, Wisconsin.

423706088363400 DELAVAN LAKE NEAR DELAVAN, WI

LOCATION.--Lat 42°36'27", long 88°36'19", in SW 1/4 NE 1/4 sec.28, T.2 N., R.16 E., Walworth County, Hydrologic Unit 07090001, at Delavan Lake Sanitary District Lift Station No. 2 at Delavan Lake Yacht Club, 1.0 mi southeast of outlet, and 2.7 mi southeast of Delavan.

DRAINAGE AREA.--41.4 mi², of which 2.3 mi² is non-contributing. Area of Delavan Lake, 2,072 acres.

PERIOD OF RECORD.--October 1983 to current year. October 1983 to September 1985 data published in Water Resources Investigation series report "Water Quality and Hydrology of Delavan Lake in Southeastern Wisconsin" by S. J. Field and M. D. Duerk (1988).

GAGE.--Water-stage recorder. Datum of gage is 922.92 ft above sea level. Prior to Sept. 5, 1989, staff gage at bridge on North Shore Drive at same datum.


REMARKS.--Lake was ice covered from Jan. 9 to Mar. 24. Lake levels controlled by Delavan Lake Sanitary District. Gage-height telemeter at station.

EXTREMES FOR PERIOD OF RECORD.--Maximum gage height observed, 6.19 ft, Feb. 21, 1994; minimum daily, -4.44 ft, Nov. 6, 1989 (lake drawn down for lake rehabilitation program).

EXTREMES FOR CURRENT YEAR.--Maximum gage height, 5.69 ft, May 23; minimum, 4.15 ft, Oct. 24.

GAGE HEIGHT, FEET WATER YEAR OCTOBER 2003 TO SEPTEMBER 2004 DAILY MEAN VALUES												
DAY	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP
1	4.26	4.17	4.78	5.00	4.96	4.93	4.97	4.98	5.11	4.99	4.85	4.82
2	4.24	4.28	4.77	5.00	4.96	4.95	4.96	4.98	5.04	4.99	4.85	4.80
3	4.23	4.42	4.77	5.00	4.98	4.95	4.96	4.98	5.00	4.99	4.86	4.79
4	4.23	4.57	4.77	5.00	4.98	4.95	4.96	4.97	5.01	5.05	4.93	4.79
5	4.22	4.66	4.79	4.99	4.98	5.07	4.95	4.97	5.03	5.03	4.91	4.78
6	4.21	4.68	4.78	4.98	5.00	5.03	4.94	4.98	5.05	5.00	4.89	4.78
7	4.20	4.67	4.79	4.97	5.00	4.96	4.95	4.97	5.06	5.01	4.88	4.77
8	4.20	4.67	4.79	4.95	5.00	4.95	4.96	4.99	5.06	5.00	4.88	4.76
9	4.19	4.66	4.81	4.95	4.99	4.95	4.96	5.03	5.06	4.99	4.87	4.75
10	4.19	4.66	5.00	4.94	4.99	4.94	4.96	5.05	5.08	4.98	4.85	4.75
11	4.19	4.66	5.09	4.94	4.99	4.94	4.96	5.05	5.11	4.96	4.84	4.74
12	4.21	4.66	5.11	4.93	4.99	4.94	4.96	5.04	5.09	4.97	4.82	4.73
13	4.20	4.65	5.11	4.93	4.99	4.93	4.95	5.11	5.02	4.97	4.81	4.73
14	4.23	4.65	5.12	4.94	4.98	4.94	4.95	5.23	4.97	4.97	4.80	4.72
15	4.23	4.64	5.12	4.93	4.98	4.94	4.95	5.29	5.01	4.97	4.79	4.72
16	4.21	4.64	5.11	4.93	4.98	4.93	4.95	5.24	5.03	4.98	4.78	4.72
17	4.20	4.65	5.11	4.96	4.97	4.93	4.98	5.18	5.12	4.96	4.78	4.71
18	4.19	4.72	e5.10	4.97	4.97	4.94	4.98	5.16	5.17	4.95	4.78	4.70
19	4.18	4.76	5.09	4.96	4.97	4.94	4.98	5.13	5.09	4.94	4.77	4.69
20	4.18	4.76	5.08	4.96	4.98	4.94	5.00	5.15	5.00	4.93	4.76	4.68
21	4.18	4.77	5.06	4.96	5.00	4.94	5.07	5.18	4.98	4.95	4.75	4.66
22	4.17	4.77	5.05	4.96	5.01	4.93	5.04	5.46	5.03	4.98	4.73	4.65
23	4.16	4.79	5.04	4.96	5.03	4.93	5.00	5.66	5.00	4.98	4.73	4.64
24	4.17	4.79	5.03	4.96	5.04	4.97	4.98	5.65	5.00	4.96	4.74	4.63
25	4.20	4.79	5.02	4.96	5.03	5.00	4.98	5.54	5.01	4.94	4.76	4.62
26	4.19	4.78	5.01	4.96	5.01	5.10	4.98	5.39	5.02	4.92	4.76	4.61
27	4.18	4.79	5.00	4.97	4.97	5.12	4.97	5.27	5.03	4.91	4.78	4.61
28	4.18	4.79	5.02	4.97	4.94	5.11	4.94	5.14	5.03	4.90	4.82	4.59
29	4.17	4.79	5.03	4.97	4.92	5.14	4.94	5.05	5.02	4.89	4.83	4.58
30	4.17	4.78	5.02	4.97	---	5.09	4.96	5.02	5.01	4.88	4.83	4.58
31	4.18	---	5.01	4.97	---	5.03	---	5.12	---	4.87	4.83	---
MEAN	4.20	4.67	4.98	4.96	4.99	4.98	4.97	5.16	5.04	4.96	4.81	4.70
MAX	4.26	4.79	5.12	5.00	5.04	5.14	5.07	5.66	5.17	5.05	4.93	4.82
MIN	4.16	4.17	4.77	4.93	4.92	4.93	4.94	4.97	4.97	4.87	4.73	4.58

e Estimated



423556088365001 DELAVAN LAKE AT CENTER NEAR DELAVAN LAKE, WI

LOCATION.--Lat 42°35'56", long 88°36'50", in SE 1/4 SW 1/4 sec.28, T.2 N., R.16 E., Walworth County, Hydrologic Unit 07090001, 2.6 mi southeast of Delavan.


DRAINAGE AREA.--41.4 mi², of which 2.3 mi² is non-contributing. Area of Delavan Lake, 2,072 acres.

PERIOD OF RECORD.--October 1983 to current year.

REMARKS.--Lake ice-covered during February measurements. Water-quality analyses done by the U.S. Geological Survey National Water Quality Laboratory. Samples for determination of chlorophyll *a* concentration are collected from the top 0.5 m of the lake.

WATER-QUALITY DATA, NOVEMBER 17, 2003 TO MAY 19, 2004
(Milligrams per liter unless otherwise indicated)

Date	Nov-17	Feb-25	Apr-13	May-19
Lake stage (ft)	4.65	5.03	4.93	5.13
Secchi depth (m)	3.0	4.9	3.4	5.0
Depth of sample (m)	0.5	16	0.5	0.5
Chlorophyll <i>a</i> , phytoplankton ($\mu\text{g/L}$)	3.4	--	6.2	--
Water temperature (°C)	7.7	7.4	0.7	7.4
Specific conductance ($\mu\text{S/cm}$)	562	564	571	566
pH	8.1	8.1	8.8	8.4
Dissolved oxygen (mg/L)	10.4	9.3	15.1	14.5
Phosphorus, total (as P)	0.102	0.116	0.086	0.087
Phosphorus, ortho, dissolved (as P)	0.074	0.074	0.052	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	0.072	--	0.206	--
Nitrogen, ammonia, dissolved (as N)	0.133	--	0.051	--
Nitrogen, amm. + org., total (as N)	0.78	--	0.66	--
Nitrogen, total (as N)	0.85	--	0.87	--
Color (Pt-Co. scale)	--	--	--	10
Turbidity (NTU)	--	--	--	<2.0
Hardness, as CaCO_3	--	--	--	230
Calcium, dissolved (Ca)	--	--	--	34.9
Magnesium, dissolved (Mg)	--	--	--	33.6
Sodium, dissolved (Na)	--	--	--	27
Potassium, dissolved (K)	--	--	--	2.73
Alkalinity, as CaCO_3	--	--	--	182
Sulfate, dissolved (SO_4)	--	--	--	23.2
Chloride, dissolved (Cl)	--	--	--	58.7
Silica, dissolved (SiO_2)	--	--	--	0.1
Solids, dissolved, at 180°C	--	--	--	335
Iron, dissolved (Fe) ($\mu\text{g/L}$)	--	--	--	<6
Manganese, dissolved, (Mn) ($\mu\text{g/L}$)	--	--	--	1
				0.9



423556088365001 DELAVAN LAKE AT CENTER NEAR DELAVAN LAKE, WI--CONTINUED


WATER-QUALITY DATA, JUNE 22 TO JULY 20, 2004
 (Milligrams per liter unless otherwise indicated)

Date	Jun-22					Jul-20		
Lake stage (ft)	5.03					4.93		
Secchi depth (m)	2.2					1.4		
Depth of sample (m)	0.5	8	12	16	0.5	6	13	16
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	30.3	--	--	--	10.4	--	--	--
Water temperature ($^{\circ}\text{C}$)	21.5	20.3	15.9	14.3	24.7	23.2	15	14.3
Specific conductance ($\mu\text{S/cm}$)	543	552	573	591	513	526	591	608
pH	8.5	8.3	8.1	8.0	9.0	8.9	9.0	8.9
Dissolved oxygen (mg/L)	10.1	5.8	1.9	0.2	11.2	7.6	0.2	0.2
Phosphorus, total (as P)	0.063	0.068	0.132	0.260	0.031	0.026	0.360	0.470
Phosphorus, ortho, dissolved (as P)	0.005	0.034	0.099	0.213	<0.006	0.003	0.305	0.387
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	0.097	--	--	--	<0.016	--	--	--
Nitrogen, ammonia, dissolved (as N)	0.042	--	--	--	0.017	--	--	--
Nitrogen, amm. + org., total (as N)	0.96	--	--	--	0.82	--	--	--
Nitrogen, total (as N)	1.1	--	--	--	--	--	--	--


6-22-04

7-20-04

DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER



PH IN STANDARD UNITS



SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS


423556088365001 DELAVAN LAKE AT CENTER NEAR DELAVAN LAKE, WI--CONTINUED

WATER-QUALITY DATA, AUGUST 19, 2004
 (Milligrams per liter unless otherwise indicated)

Date	<u>Aug-19</u>							
Lake stage (ft)	4.77							
Secchi depth (m)	1.5							
Depth of sample (m)	0.5	5	9	11	13	14	15	16
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	20.6	--	--	--	--	--	--	--
Water temperature ($^{\circ}\text{C}$)	21.8	20.7	20.6	18.6	16.5	14.8	14.6	14.4
Specific conductance ($\mu\text{S/cm}$)	526	529	533	563	587	607	611	620
pH	8.6	8.4	8.4	7.9	7.7	7.7	7.7	7.7
Dissolved oxygen (mg/L)	9.6	7.5	6.5	0.2	0.2	0.2	0.2	0.2
Phosphorus, total (as P)	0.088	0.072	0.069	0.155	0.360	0.490	0.520	0.560
Phosphorus, ortho, dissolved (as P)	0.012	--	0.028	--	--	0.459	--	0.554
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	0.008	--	--	--	--	--	--	--
Nitrogen, ammonia, dissolved (as N)	0.042	--	--	--	--	--	--	--
Nitrogen, amm. + org., total (as N)	0.82	--	--	--	--	--	--	--


8-19-04

DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER



WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS

PH IN STANDARD UNITS




SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS

423556088365001 DELAVAN LAKE AT CENTER NEAR DELAVAN LAKE, WI--CONTINUED


WATER-QUALITY DATA, SEPTEMBER 22, 2004
 (Milligrams per liter unless otherwise indicated)

Date	<u>Sep-22</u>			
Lake stage (ft)	4.68			
Secchi depth (m)	1.7			
Depth of sample (m)	0.5	12	14	16
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	25.0	--	--	--
Water temperature ($^{\circ}\text{C}$)	21.2	20.0	16.4	14.9
Specific conductance ($\mu\text{S/cm}$)	525	545	600	636
pH	8.8	8.2	7.9	7.8
Dissolved oxygen (mg/L)	8.6	0.7	0.2	0.2
Phosphorus, total (as P)	0.107	0.156	0.380	0.710
Phosphorus, ortho, dissolved (as P)	0.044	0.097	0.355	0.683
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	0.011	--	--	--
Nitrogen, ammonia, dissolved (as N)	0.041	--	--	--
Nitrogen, amm. + org., total (as N)	0.94	--	--	--


9-22-04
DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER



PH IN STANDARD UNITS



SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS



423556088365001 DELAVAN LAKE AT CENTER NEAR DELAVAN LAKE, WI--CONTINUED

ADDITIONAL WATER-QUALITY DATA, OCTOBER 1, 2003 TO SEPTEMBER 30, 2004*
 (Milligrams per liter unless otherwise indicated)

	<u>Oct. 1</u>	<u>Oct. 8</u>	<u>Oct. 17</u>	<u>Oct. 22</u>	<u>Oct. 30</u>
Lake stage (ft)	4.26	4.20	4.20	4.17	4.17
Secchi-depth (meters)	1.5	2.0	2.4	3.0	3.2
Depth of sample (meters)	0.5	0.5	0.5	0.5	0.5
Water temperature (°C)	15.0	14.0	14.0	13.5	12.0
Phosphorus, total (as P)	0.116	0.121	0.118	0.109	0.108

	<u>Nov. 6</u>	<u>Nov. 14</u>	<u>April 1</u>	<u>April 7</u>	<u>April 22</u>
Lake stage (ft)	4.68	4.65	4.97	4.95	5.04
Secchi-depth (meters)	3.0	3.0	3.5	3.7	5.3
Depth of sample (meters)	0.5	0.5	0.5	0.5	0.5
Water temperature (°C)	10.0	7.5	6.5	8.0	11.0
Phosphorus, total (as P)	0.103	0.105	0.068	0.059	0.067

	<u>April 27</u>	<u>May 4</u>	<u>May 11</u>	<u>May 26</u>	<u>June 4</u>
Lake stage (ft)	4.97	4.97	5.05	5.39	5.01
Secchi-depth (meters)	6.4	6.1	4.9	3.4	4.4
Depth of sample (meters)	0.5	0.5	0.5	0.5	0.5
Water temperature (°C)	11.0	12.0	15.0	17.8	18.0
Phosphorus, total (as P)	0.051	0.050	0.051	0.054	0.066

	<u>June 9</u>	<u>June 18</u>	<u>June 22</u>	<u>July 1</u>	<u>July 9</u>
Lake stage (ft)	5.06	5.17	5.03	4.99	4.99
Secchi-depth (meters)	3.0	1.2	2.6	2.0	1.5
Depth of sample (meters)	0.5	0.5	0.5	0.5	0.5
Water temperature (°C)	22.0	24.5	21.5	23.0	22.0
Phosphorus, total (as P)	0.051	0.182	0.056	0.065	0.074

	<u>July 15</u>	<u>July 28</u>	<u>Aug. 2</u>	<u>Aug. 13</u>	<u>Aug. 18</u>
Lake stage (ft)	4.97	4.90	4.85	4.81	4.78
Secchi-depth (meters)	2.0	1.5	1.8	1.2	1.7
Depth of sample (meters)	0.5	0.5	0.5	0.5	0.5
Water temperature (°C)	24.0	24.0	24.0	21.0	21.5
Phosphorus, total (as P)	0.026	0.048	0.050	0.148	0.077

	<u>Aug. 27</u>	<u>Aug. 31</u>	<u>Sept. 9</u>	<u>Sept. 17</u>	<u>Sept. 30</u>
Lake stage (ft)	4.78	4.83	4.75	4.71	4.58
Secchi-depth (meters)	2.4	2.4	1.5	1.7	1.7
Depth of sample (meters)	0.5	0.5	0.5	0.5	0.5
Water temperature (°C)	22.0	23.0	23.0	21.0	22.8
Phosphorus, total (as P)	0.099	0.111	0.095	0.103	0.100

* Measurements and samples collected by the Delavan Lake Sanitary District.

423659088354401 DELAVAN LAKE, AT NORTH END, NEAR LAKE LAWN, WI

LOCATION.--Lat 42°36'59", long 88°35'44", in NW 1/4 SW 1/4, sec.22, T.2 N., R.16 E., Walworth County, Hydrologic Unit 07090001, 2.6 mi southeast of Delavan.

DRAINAGE AREA.--41.4 mi², of which 2.3 mi² is non-contributing.

PERIOD OF RECORD.--October 1983 to current year.

WATER-QUALITY DATA, APRIL 13 TO AUGUST 19, 2004

	April 13	May 19	June 22	July 20	Aug. 19
Secchi depth (meters)	3.4	5.3	2.2	1.3	1.3

423526088380101 DELAVAN LAKE, AT SW END, NEAR DELAVAN LAKE, WI

LOCATION.--Lat 42°35'26", long 88°38'01", in SE 1/4 NW 1/4, sec.32, T.2 N., R.16 E., Walworth County, Hydrologic Unit 07090001, 2.6 mi southeast of Delavan.

DRAINAGE AREA.--41.4 mi², of which 2.3 mi² is non-contributing.

PERIOD OF RECORD.--October 1983 to current year.

WATER-QUALITY DATA, APRIL 13 TO AUGUST 19, 2004

	April 13	May 19	June 22	July 20	Aug. 19
Secchi depth (meters)	3.0	3.8	2.2	1.4	1.7

453446091465100 LAKE DESAIR NEAR RICE LAKE, WI

LOCATION.--Lat 45°34'46", long 91°46'51", in SE 1/4 SW 1/4 NW 1/4 sec.6, T.35 N., R.11 W., Barron County, Hydrologic Unit 07050007, at public boat landing off 18th Street about 3 miles northwest of the city of Rice Lake.


PERIOD OF RECORD.--August 2004 (discontinued). Data collected through Wis. Dept. of Natural Resources Self-Help Monitoring program since April 1993 are available.

REMARKS.--Lake sampled at deepest location. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, AUGUST 17, 2004
(Milligrams per liter unless otherwise indicated)


Date	Aug-17				
Secchi depth (m)		0.85			
Depth of sample (m)	0.5	3	3.5	6	8
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	91.7	--	--	--	--
Water temperature ($^{\circ}\text{C}$)	19.3	18.8	17.6	10.5	9.3
Specific conductance ($\mu\text{S/cm}$)	65	67	69	122	147
pH	6.2	7.3	7.1	6.9	6.8
Dissolved oxygen (mg/L)	9.6	4.4	0.5	0.1	0.1
Phosphorus, total (as P)	0.058	0.058	0.054	0.524	0.922

8-17-04
DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER



WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS

PH IN STANDARD UNITS



SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS

05404500 DEVILS LAKE NEAR BARABOO, WI

LOCATION.--Lat 43°25'35", long 89°43'40", in SW 1/4 SE 1/4 sec.13, T.11 N., R.6 E., Sauk County, Hydrologic Unit 07070004, in Devils Lake State Park, 3.5 mi south of Baraboo.

DRAINAGE AREA.--4.79 mi². Area of Devils Lake, 361 acres.

PERIOD OF RECORD.--June 1922 to August 1930, June to August 1932, June 1934 to September 1981, October 1984 to June 1991 (fragmentary), July 1991 to current year. Unpublished daily stage records from October 1981 to September 1984 in District files.

REVISED RECORDS.--WDR WI-78-1: Drainage area.

GAGE.--Water-stage recorder installed July 17, 1991. Datum of gage is 955.00 ft, above sea level.


REMARKS.--Lake has no surface outlet. Water removed from lake by pumping or siphon Oct. 1-15 and Sept. 8-30.

EXTREMES FOR PERIOD OF RECORD.--Maximum gage height observed, 14.13 ft, July 18, 1993; minimum observed, 1.49 ft, Feb. 8, 1965.

EXTREMES FOR CURRENT YEAR.--Maximum recorded gage height, 11.86 ft, June 17, 18; minimum recorded, 6.66 ft, Nov. 1.

GAGE HEIGHT, FEET
WATER YEAR OCTOBER 2003 TO SEPTEMBER 2004
DAILY MEAN VALUES

DAY	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP
1	7.08	6.66	6.95	7.13	7.13	7.27	8.92	8.86	10.67	11.24	10.77	10.10
2	7.04	6.67	6.95	7.14	7.14	7.36	8.93	8.84	10.68	11.24	10.77	10.05
3	7.00	6.73	6.94	7.15	7.15	7.42	8.93	8.83	10.68	11.21	10.79	10.00
4	6.98	6.90	6.94	7.16	7.15	7.47	8.94	8.81	10.67	11.31	10.89	9.98
5	6.95	6.90	6.94	7.16	7.15	7.79	8.94	8.79	10.66	11.32	10.86	9.96
6	6.94	6.89	6.94	7.17	7.19	8.00	8.94	8.78	10.66	11.31	10.83	9.93
7	6.90	6.88	6.93	7.16	7.19	8.08	8.95	8.76	10.66	11.27	10.80	9.89
8	6.87	6.86	6.93	7.16	7.19	8.13	8.93	8.79	10.64	11.22	10.77	9.83
9	6.86	6.85	6.92	7.16	7.19	8.17	8.93	8.91	10.68	11.19	10.74	9.78
10	6.86	6.84	7.00	7.15	7.18	8.20	8.92	8.94	11.17	11.17	10.70	9.73
11	6.85	6.83	7.02	7.15	7.18	8.25	8.90	8.96	11.50	11.13	10.66	9.68
12	6.85	6.83	7.02	7.15	7.18	8.27	8.89	8.98	11.65	11.09	10.62	9.63
13	6.84	6.81	7.01	7.15	7.18	8.28	8.88	9.04	11.72	11.04	10.59	9.59
14	6.86	6.80	7.01	7.15	7.17	8.32	8.86	9.12	11.76	10.98	10.56	9.53
15	6.84	6.80	7.01	7.15	7.17	8.34	8.85	9.14	11.76	10.93	10.54	9.52
16	6.83	6.79	7.02	7.14	7.17	8.35	8.84	9.15	11.77	10.98	10.51	9.50
17	6.82	6.80	7.02	7.15	7.16	8.37	8.85	9.16	11.86	11.11	10.49	9.45
18	6.80	6.81	7.02	7.15	7.16	8.38	8.92	9.19	11.86	11.10	10.47	9.40
19	6.79	6.81	7.01	7.14	7.16	8.39	8.92	9.19	11.81	11.08	10.44	9.34
20	6.78	6.80	7.01	7.14	7.18	8.41	8.92	9.21	11.76	11.06	10.38	9.28
21	6.77	6.78	7.00	7.13	7.20	8.42	8.94	9.30	11.71	11.07	10.32	9.22
22	6.76	6.78	7.00	7.12	7.20	8.43	8.94	9.81	11.68	11.06	10.26	9.18
23	6.75	6.96	7.00	7.13	7.21	8.43	8.93	10.11	11.63	11.03	10.23	9.13
24	6.74	7.00	6.99	7.13	7.21	8.46	8.92	10.28	11.61	11.00	10.22	9.08
25	6.73	7.00	6.99	7.13	7.21	8.50	8.93	10.34	11.56	10.97	10.25	9.03
26	6.71	6.99	6.99	7.14	7.21	8.64	8.92	10.35	11.50	10.94	10.20	8.98
27	6.70	6.99	6.99	7.15	7.21	8.69	8.91	10.36	11.45	10.92	10.32	8.94
28	6.69	6.98	7.07	7.14	7.20	8.75	8.90	10.35	11.40	10.89	10.28	8.88
29	6.68	6.98	7.11	7.14	7.21	8.82	8.89	10.39	11.35	10.85	10.24	8.83
30	6.68	6.97	7.12	7.14	---	8.87	8.87	10.47	11.29	10.83	10.19	8.79
31	6.67	---	7.13	7.13	---	8.90	--	10.62	---	10.80	10.14	---
MEAN	6.83	6.86	7.00	7.14	7.18	8.26	8.91	9.41	11.33	11.08	10.51	9.47
MAX	7.08	7.00	7.13	7.17	7.21	8.90	8.95	10.62	11.86	11.32	10.89	10.10
MIN	6.67	6.66	6.92	7.12	7.13	7.27	8.84	8.76	10.64	10.80	10.14	8.79



05406050 FISH LAKE NEAR SAUK CITY, WI

LOCATION.--Lat 43°17'27", long 89°39'09" in NE 1/4 SW 1/4 sec.3, T.9 N., R.7 E., Dane County, Hydrologic Unit 07070005, on north side of lake, 0.4 mi southwest of Crystal Lake, and 3.1 mi east of Sauk City.

DRAINAGE AREA.--2.23 mi². Area of Fish Lake, 252 acres.

PERIOD OF RECORD.--November 1966 to September 1981, April 1985 to May 1987, May 1988, April 1989 to October 1990 (fragmentary); continuous record from October 1990 to November 1996; nonrecording gage November 1996 to current year.

REVISED RECORDS.--WDR WI-92-1: Drainage area. WDR WI-87-1: All published values for the 1987 water year are invalid. Two valid values for water years 1987 and 1988 are available: May 7, 1987, water surface 10.52 ft, and May 16, 1988, water surface 10.83 ft.

GAGE.--Nonrecording gage. Datum of gage is 848.07 ft above sea level. Local observer, Richard Lillie, reads staff gage when lake is ice-free.


REMARKS.--Lake has no surface outlet.

EXTREMES FOR PERIOD OF RECORD.--Maximum gage height observed, 13.39 ft, June 14 and 26, 2002; minimum observed, 3.02 ft, Aug. 29, 1970.

EXTREMES FOR CURRENT YEAR.--Maximum gage height observed, 12.07 ft, Mar. 23; minimum observed, 11.33 ft, Oct. 30.

GAGE HEIGHT, FEET
WATER YEAR OCTOBER 2003 TO MARCH 2004
DAILY MEAN VALUES

DAY	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP
1	---	---	---	---	---	---	---	---	---	---	---	---
2	11.50	11.36	11.77	---	---	---	---	---	---	---	---	---
3	11.48	11.46	---	---	---	---	---	---	---	---	---	---
4	---	11.78	---	---	---	---	---	---	---	---	---	---
5	11.47	11.78	---	---	---	---	---	---	---	---	---	---
6	---	---	---	---	---	---	---	---	---	---	---	---
7	---	---	---	---	---	---	---	---	---	---	---	---
8	11.45	11.75	---	---	---	---	---	---	---	---	---	---
9	---	---	---	---	---	---	---	---	---	---	---	---
10	---	---	---	---	---	---	---	---	---	---	---	---
11	11.44	---	---	---	---	---	---	---	---	---	---	---
12	11.43	11.71	---	---	---	---	---	---	---	---	---	---
13	---	---	---	---	---	---	---	---	---	---	---	---
14	11.48	---	---	---	---	---	---	---	---	---	---	---
15	---	11.69	---	---	---	---	---	---	---	---	---	---
16	---	---	---	---	---	---	---	---	---	---	---	---
17	---	---	---	---	---	---	---	---	---	---	---	---
18	---	11.71	---	---	---	---	---	---	---	---	---	---
19	11.40	---	---	---	---	---	---	---	---	---	---	---
20	---	---	---	---	---	---	---	---	---	---	---	---
21	11.39	---	---	---	---	---	---	---	---	---	---	---
22	---	11.68	---	---	---	---	---	---	---	---	---	---
23	11.38	11.80	---	---	---	12.07	---	---	---	---	---	---
24	---	---	---	---	---	---	---	---	---	---	---	---
25	11.37	---	---	---	---	---	---	---	---	---	---	---
26	---	---	---	---	---	---	---	---	---	---	---	---
27	---	---	---	---	---	---	---	---	---	---	---	---
28	---	---	---	---	---	---	---	---	---	---	---	---
29	11.35	---	---	---	---	---	---	---	---	---	---	---
30	11.33	11.80	---	---	---	---	---	---	---	---	---	---
31	---	---	---	---	---	---	---	---	---	---	---	---
MEAN	---	---	---	---	---	---	---	---	---	---	---	---
MAX	---	---	---	---	---	---	---	---	---	---	---	---
MIN	---	---	---	---	---	---	---	---	---	---	---	---



433632088100200 FOREST LAKE NEAR DUNDEE, WI

LOCATION.--Lat 43°36'32", long 88°10'02", in SW 1/4 NE 1/4 sec.12, T.13 N., R.19 E., Fond du Lac County, Hydrologic Unit 04040003, 3 mi south of Dundee.

PERIOD OF RECORD.--March 1994 to August 1996, May to August 2004.

REMARKS.--Lake sampled near center at the deep hole. Lake ice-covered during February sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, MAY 3 TO AUGUST 18, 2004
(Milligrams per liter unless otherwise indicated)

Date	May-3	Jun-9	Jul-14	Aug-18
Secchi depth (m)	1.9	5.0	3.3	3.6
Depth of sample (m)	0.5	8.5	0.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	10.6	--	4.6	--
Water temperature (°C)	12.6	8.1	25.1	9.7
Specific conductance ($\mu\text{S/cm}$)	249	259	230	247
pH	8.4	7.7	8.3	7.1
Dissolved oxygen (mg/L)	11.4	3.0	11.4	0.5
Phosphorus, total (as P)	0.020	0.041	0.018	0.082
Phosphorus, ortho, dissolved (as P)	0.003	--	--	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	<0.019	--	--	--
Nitrogen, ammonia, dissolved (as N)	<0.015	--	--	--
Nitrogen, amm. + org., total (as N)	0.81	--	--	--
Color (Pt-Co. scale)	5	--	--	--
Turbidity (NTU)	1.9	--	--	--
Hardness, as CaCO_3	116	--	--	--
Calcium, dissolved (Ca)	26.6	--	--	--
Magnesium, dissolved (Mg)	14.5	--	--	--
Sodium, dissolved (Na)	2.5	--	--	--
Potassium, dissolved (K)	<1	--	--	--
Alkalinity, as CaCO_3	116	--	--	--
Sulfate, dissolved (SO_4)	<4.5	--	--	--
Chloride, dissolved (Cl)	4.9	--	--	--
Silica, dissolved (SiO_2)	0.126	--	--	--
Solids, dissolved, at 180°C	128	--	--	--
Iron, dissolved (Fe) ($\mu\text{g/L}$)	<100	--	--	--
Manganese, dissolved, (Mn) ($\mu\text{g/L}$)	1.0	--	--	--





5-3-04

6-9-04

7-14-04





8-18-04

DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER




WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS

PH IN STANDARD UNITS



SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS



423525088260400 GENEVA LAKE AT LAKE GENEVA, WI

LOCATION.--Lat 42°35'25", long 88°26'04" in SE 1/4 NW 1/4 sec.36, T.2 N., R.17 E., Walworth County, Hydrologic Unit 07120006, at Geneva Lake dam at Center Street at Lake Geneva.

DRAINAGE AREA.--28.7 mi². Area of Geneva Lake, 5,262 acres.

PERIOD OF RECORD.--October 1997 to August 2002, December 2002 to current year.

GAGE.--Water-stage recorder. Datum of gage is 862.08 ft above sea level. Intermittent staff-gage readings January to February.


REMARKS.--Gage-height telemeter at station.

EXTREMES FOR PERIOD OF RECORD.--Maximum gage height, 3.29 ft, June 13, 2000; minimum gage height, 1.50 ft, Oct. 11, 2003 (affected by wind).

EXTREMES FOR CURRENT YEAR.--Maximum gage height, 3.28 ft, May 28; minimum gage height, 1.50 ft, Oct. 11 (affected by wind).

GAGE HEIGHT, FEET WATER YEAR OCTOBER 2003 TO SEPTEMBER 2004 DAILY MEAN VALUES													
DAY	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	
1	1.75	1.67	1.99	2.19	---	2.46	2.56	2.51	2.92	2.59	2.36	2.27	
2	1.73	1.77	1.97	2.20	---	2.48	2.55	2.50	2.87	2.56	2.35	2.27	
3	1.74	1.87	1.96	2.21	---	2.48	2.54	2.49	2.83	2.56	2.34	2.27	
4	1.71	1.96	1.95	2.20	e2.27	2.49	2.53	2.50	2.80	2.62	2.39	2.27	
5	1.70	2.02	1.97	2.22	---	2.62	2.52	2.48	2.77	2.59	2.37	2.28	
6	1.69	2.02	1.97	---	---	2.60	2.52	2.48	2.75	2.60	2.36	2.29	
7	1.69	2.01	1.97	---	---	2.59	2.51	2.45	2.74	2.58	2.35	2.26	
8	1.70	1.98	1.97	---	---	2.56	2.51	2.48	2.73	2.55	2.34	2.23	
9	1.69	1.98	1.98	---	---	2.53	2.50	2.50	2.71	2.53	2.34	2.22	
10	1.69	1.98	2.15	---	---	2.51	2.49	2.51	2.71	2.54	2.32	2.22	
11	1.70	1.98	2.19	---	---	2.50	2.48	2.51	2.74	2.54	2.29	2.21	
12	1.71	2.01	2.18	---	---	2.47	2.47	2.54	2.78	2.54	2.26	2.21	
13	1.70	1.97	2.17	---	---	2.44	2.46	2.71	2.78	2.54	2.24	2.21	
14	1.74	1.96	2.18	---	---	2.46	2.46	2.79	2.76	2.52	2.23	2.20	
15	1.74	1.94	2.19	---	---	2.42	2.46	2.81	2.74	2.51	2.23	2.22	
16	1.72	1.94	2.21	---	---	2.40	2.46	2.79	2.71	2.50	2.23	2.20	
17	1.71	1.94	2.19	---	---	2.40	2.47	2.78	2.75	2.47	2.22	2.17	
18	1.71	2.02	2.18	---	---	2.39	2.50	2.84	2.76	2.47	2.23	2.16	
19	1.69	2.05	2.18	---	---	2.38	2.51	2.81	2.71	2.47	2.21	2.16	
20	1.71	2.05	2.18	---	---	2.38	2.48	2.81	2.69	2.46	2.19	2.15	
21	1.69	2.02	2.19	---	---	2.35	2.56	2.85	2.70	2.47	2.18	2.14	
22	1.68	2.02	2.17	---	---	2.34	2.53	3.01	2.73	2.49	2.19	2.12	
23	1.67	2.07	2.17	---	---	2.32	2.53	3.06	2.71	2.46	2.16	2.12	
24	1.67	2.09	2.17	---	---	2.36	2.51	3.04	2.71	2.44	2.17	2.11	
25	1.70	2.04	2.17	---	e2.42	2.38	2.55	2.99	2.70	2.42	2.21	2.08	
26	1.70	2.01	2.16	---	---	2.48	2.55	2.95	2.68	2.41	2.22	2.08	
27	1.68	2.01	2.16	---	2.42	2.49	2.53	2.91	2.65	2.40	2.24	2.07	
28	1.70	2.02	2.20	e2.28	2.42	2.53	2.54	2.85	2.64	2.39	2.25	2.04	
29	1.67	2.02	2.21	---	2.45	2.58	2.53	2.82	2.62	2.38	2.28	2.03	
30	1.67	2.00	2.21	---	---	2.57	2.50	2.87	2.61	2.37	2.28	2.02	
31	1.68	---	2.21	---	---	2.57	---	2.95	---	2.36	2.27	---	
MEAN	1.70	1.98	2.12	---	---	2.47	2.51	2.73	2.73	2.49	2.27	2.18	
MAX	1.75	2.09	2.21	---	---	2.62	2.56	3.06	2.92	2.62	2.39	2.29	
MIN	1.67	1.67	1.95	---	---	2.32	2.46	2.45	2.61	2.36	2.16	2.02	

e Estimated



423329088323300 GENEVA LAKE AT WEST END NEAR WILLIAMS BAY, WI

LOCATION.--Lat 42°33'29", long 88°32'33", in NE 1/4 SE 1/4, sec.12, T.1 N., R.16 E., Walworth County, Hydrologic Unit 07120006, 1.3 mi south of Williams Bay.

DRAINAGE AREA.--28.7 mi².

PERIOD OF RECORD.--April 1997 to current year.


REMARKS.--Lake sampled at deep hole at a depth of about 43 m. Water-quality analyses done by Wisconsin State Laboratory of Hygiene. Samples for determination of chlorophyll *a* concentration are collected from the top 0.5 m of the lake.

WATER-QUALITY DATA, OCTOBER 24, 2003 TO APRIL 13, 2004
(Milligrams per liter unless otherwise indicated)

Date	<u>Oct-24</u>		<u>Apr-13</u>	
Lake stage (ft)	1.67		2.46	
Secchi depth (m)	6.2		5.3	
Depth of sample (m)	0.5	18	27	42
Chlorophyll <i>a</i> , phytoplankton (µg/L)	3.54	--	--	3.86
Water temperature (°C)	13.5	13.4	8.7	7.9
Specific conductance (µS/cm)	523	523	530	532
pH	8.1	8.1	7.6	7.4
Dissolved oxygen (mg/L)	9.6	8.1	3.6	0.2
Phosphorus, total (as P)	0.013	0.013	0.015	0.055
Phosphorus, ortho, dissolved (as P)	0.003	--	--	0.002
Nitrogen, NO ₂ + NO ₃ , diss. (as N)	<0.022	--	--	0.048
Nitrogen, ammonia, dissolved (as N)	<0.013	--	--	0.025
Nitrogen, amm. + org., total (as N)	0.32	--	--	0.38
Nitrogen, total (as N)	--	--	--	0.43
Color (Pt-Co. scale)	--	--	--	5
Turbidity (NTU)	--	--	--	<1
Hardness, as CaCO ₃	--	--	--	230
Calcium, dissolved (Ca)	--	--	--	34.4
Magnesium, dissolved (Mg)	--	--	--	34.5
Sodium, dissolved (Na)	--	--	--	18.4
Potassium, dissolved (K)	--	--	--	2
Alkalinity, as CaCO ₃	--	--	--	184
Sulfate, dissolved (SO ₄)	--	--	--	31.1
Chloride, dissolved (Cl)	--	--	--	37.5
Silica, dissolved (SiO ₂)	--	--	--	1.73
Solids, dissolved, at 180°C	--	--	--	282
Iron, dissolved (Fe) (µg/L)	--	--	--	<100
Lead, (Pb) (µg/L)	--	--	--	<1
Manganese, dissolved, (Mn) (µg/L)	--	--	--	<1


10-24-03 4-13-04

DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER



WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS

pH IN STANDARD UNITS




SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS

423329088323300 GENEVA LAKE AT WEST END NEAR WILLIAMS BAY, WI--CONTINUED


WATER-QUALITY DATA, JUNE 22, 2004
(Milligrams per liter unless otherwise indicated)

			<u>Jun-22</u>			
Date			2.73			
Lake stage (ft)			5.3			
Secchi depth (m)						
Depth of sample (m)	0.5	8	20	33	38	42
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	4.02	--	--	--	--	--
Water temperature ($^{\circ}\text{C}$)	20	19.8	9.9	8.6	8.3	8.3
Specific conductance ($\mu\text{S/cm}$)	510	511	531	534	536	537
pH	8.3	8.3	8.7	8.4	8.2	8.1
Dissolved oxygen (mg/L)	9.7	9.6	9.1	9.3	7.8	7
Phosphorus, total (as P)	0.007	0.009	0.007	0.007	0.008	0.011
Phosphorus, ortho, dissolved (as P)	<0.002	--	--	--	--	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	<0.019	--	--	--	--	--
Nitrogen, ammonia, dissolved (as N)	<0.015	--	--	--	--	--
Nitrogen, amm. + org., total (as N)	0.3	--	--	--	--	--

6-22-04
DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER



WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS



SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS


423329088323300 GENEVA LAKE AT WEST END NEAR WILLIAMS BAY, WI--CONTINUED

WATER-QUALITY DATA, JULY 20, 2004
 (Milligrams per liter unless otherwise indicated)

			Jul-20			
Date			2.46			
Lake stage (ft)			5.3			
Secchi depth (m)						
Depth of sample (m)	0.5	8	25	33	38	42
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	1.6	--	--	--	--	--
Water temperature ($^{\circ}\text{C}$)	23	22.5	9.5	8.5	8.3	8.3
Specific conductance ($\mu\text{S/cm}$)	513	513	536	537	538	540
pH	8.4	8.4	9.1	8.6	8.2	8
Dissolved oxygen (mg/L)	9.2	9	8.3	7.7	4.8	4.1
Phosphorus, total (as P)	0.008	0.01	0.006	0.009	0.028	0.038
Phosphorus, ortho, dissolved (as P)	0.002	--	--	--	--	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	<0.019	--	--	--	--	--
Nitrogen, ammonia, dissolved (as N)	<0.015	--	--	--	--	--
Nitrogen, amm. + org., total (as N)	0.35	--	--	--	--	--


7-20-04

DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER



WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS

PH IN STANDARD UNITS



SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS


423329088323300 GENEVA LAKE AT WEST END NEAR WILLIAMS BAY, WI--CONTINUED

WATER-QUALITY DATA, AUGUST 19, 2004
(Milligrams per liter unless otherwise indicated)


Date	Aug-19					
Lake stage (ft)	2.21					
Secchi depth (m)	5.8					
Depth of sample (m)	0.5	12	21	33	38	42
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	2.65	--	--	--	--	--
Water temperature ($^{\circ}\text{C}$)	20.8	20.6	10.6	8.6	8.3	8.2
Specific conductance ($\mu\text{S/cm}$)	510	511	532	533	534	537
pH	8	8.2	8.3	8.1	7.8	7.6
Dissolved oxygen (mg/L)	8.3	7.8	6.2	4.9	1.7	0.2
Phosphorus, total (as P)	0.009	0.01	0.008	0.008	0.04	0.078
Phosphorus, ortho, dissolved (as P)	0.002	--	--	--	--	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	<0.019	--	--	--	--	--
Nitrogen, ammonia, dissolved (as N)	0.038	--	--	--	--	--
Nitrogen, amm. + org., total (as N)	0.43	--	--	--	--	--

8-19-04

DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER



WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS




SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS

423329088323300 GENEVA LAKE AT WEST END NEAR WILLIAMS BAY, WI--CONTINUED


WATER-QUALITY DATA, SEPTEMBER 22, 2004
(Milligrams per liter unless otherwise indicated)

Date	<u>Sep-22</u>					
Lake stage (ft)	2.12					
Secchi depth (m)	5.5					
Depth of sample (m)	0.5	12	25	33	38	42
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	2.61	--	--	--	--	--
Water temperature ($^{\circ}\text{C}$)	21	20.7	9.6	8.6	8.4	8.3
Specific conductance ($\mu\text{S/cm}$)	510	511	529	530	532	536
pH	8.2	8.2	8.4	8	7.8	7.7
Dissolved oxygen (mg/L)	8.3	7	5.1	2.9	0.2	0.1
Phosphorus, total (as P)	0.011	0.013	0.008	0.015	0.055	0.072
Phosphorus, ortho, dissolved (as P)	0.003	<0.002	0.003	0.008	0.045	0.057
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	<0.019	<0.019	0.123	0.263	0.292	0.188
Nitrogen, ammonia, dissolved (as N)	<0.015	<0.015	<0.015	<0.015	<0.015	0.113
Nitrogen, amm. + org., total (as N)	0.45	0.46	0.33	0.4	0.41	0.49
Nitrogen, total (as N)	--	--	0.45	0.66	0.7	0.68


9-22-04
DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER



WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS



SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS



Surface total phosphorus, chlorophyll a concentrations, Secchi depths,
and TSI data for Lake Geneva, West End, near Williams Bay, Wisconsin.

(Circles on the first two plots indicate laboratory detection limit for selected analyses.
Actual concentrations for these particular analyses are less than the plotted circles.)

434928088553601 GREEN LAKE AT COUNTY TRUNK HIGHWAY A NEAR GREEN LAKE, WI

LOCATION.--Lat 43°49'28", long 88°55'36" in NE 1/4 SE 1/4 SE 1/4 sec.27, T.16 N., R.13 E., Green Lake County, Hydrologic Unit 04030201, on left bank at downstream side of County Trunk Highway A, 2.3 mi southeast of Green Lake.

DRAINAGE AREA.--103 mi².

PERIOD OF RECORD.--October 1993 to current year.

GAGE.--Water-stage recorder. Datum of gage is 790.00 ft above sea level.

REMARKS.--Lake level regulated by dam at outlet at Green Lake. Gage-height telemeter at station.

EXTREMES FOR PERIOD OF RECORD.--Maximum recorded gage height, 7.64 ft, June 17, 2004; minimum recorded, 5.41 ft, Jan. 17, 1995.

EXTREMES FOR CURRENT YEAR.--Maximum recorded gage height, 7.64 ft, June 17; minimum recorded, 5.73 ft, Oct. 30, Nov. 1, 2.

GAGE HEIGHT, FEET
WATER YEAR OCTOBER 2003 TO SEPTEMBER 2004
DAILY MEAN VALUES

DAY	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP
1	5.91	5.75	6.30	6.37	6.32	6.42	6.60	6.56	7.50	6.91	6.31	6.27
2	5.88	5.77	6.27	6.38	6.32	6.54	6.59	6.55	7.50	6.84	6.33	6.27
3	5.90	5.78	6.27	6.40	6.35	6.60	6.58	6.54	7.49	6.79	6.32	6.27
4	5.89	5.92	6.27	6.37	6.34	6.65	6.57	6.54	7.49	6.82	6.32	6.26
5	5.87	5.96	6.27	6.39	6.34	6.81	6.55	6.52	7.41	6.79	6.31	6.25
6	5.87	5.95	6.27	6.40	6.37	6.95	6.54	6.51	7.37	6.81	6.29	6.26
7	5.87	5.95	6.27	6.36	6.37	7.03	6.53	---	7.33	6.78	6.29	6.24
8	5.87	5.92	6.27	6.34	6.37	7.04	6.52	---	7.28	6.73	6.27	6.20
9	5.87	5.92	6.27	6.34	6.37	7.02	6.51	---	7.25	6.69	6.28	6.19
10	5.86	5.90	6.34	6.34	6.36	7.00	6.49	---	7.21	6.67	6.27	6.18
11	5.87	5.92	6.39	6.33	6.35	6.98	6.48	---	7.41	6.63	6.24	6.18
12	5.90	5.96	6.37	6.33	6.35	6.91	6.46	---	7.50	6.58	6.20	6.16
13	5.87	5.94	6.36	6.32	6.35	6.86	6.45	---	7.57	6.53	6.18	6.15
14	5.89	5.91	6.36	6.32	6.34	6.84	6.44	---	7.60	6.56	6.16	6.15
15	5.88	5.91	6.36	6.32	6.33	6.79	6.43	---	7.60	6.53	6.16	6.16
16	5.86	5.91	6.39	6.32	6.33	6.75	6.43	---	7.58	6.52	6.15	6.16
17	5.85	5.91	6.38	6.33	6.33	6.72	6.43	---	7.60	6.50	6.19	6.13
18	5.84	5.99	6.37	6.33	6.32	6.69	6.49	---	7.59	6.49	6.22	6.12
19	5.83	6.02	6.37	6.33	6.31	6.67	6.53	---	7.54	6.48	6.22	6.11
20	5.84	6.02	6.36	6.32	6.33	6.66	6.53	6.92	7.51	6.46	6.20	6.09
21	5.82	6.01	6.36	6.32	6.35	6.62	6.57	6.98	7.50	6.47	6.18	6.07
22	5.81	5.99	6.35	6.31	6.35	6.60	6.57	7.10	7.41	6.46	6.17	6.06
23	5.80	6.19	6.35	6.31	6.37	6.58	6.57	7.22	7.35	6.43	6.13	6.05
24	5.79	6.28	6.35	6.32	6.37	6.57	6.57	7.36	7.35	6.40	6.13	6.04
25	5.80	6.25	6.34	6.32	6.36	6.57	6.58	7.42	7.27	6.36	6.14	6.01
26	5.79	6.26	6.33	6.33	6.36	6.58	6.59	7.43	7.19	6.35	6.15	6.00
27	5.77	6.27	6.33	6.33	6.35	6.58	6.58	7.43	7.13	6.34	6.27	5.99
28	5.78	6.29	6.38	6.33	6.35	6.60	6.57	7.38	7.09	6.33	6.27	5.96
29	5.76	6.28	6.40	6.33	6.36	6.62	6.58	7.34	7.05	6.31	6.27	5.96
30	5.75	6.30	6.40	6.33	---	6.61	6.56	7.35	6.99	6.28	6.27	5.95
31	5.76	---	6.40	6.33	---	6.60	---	7.46	---	6.28	6.27	---
MEAN	5.84	6.01	6.34	6.34	6.35	6.72	6.53	---	7.39	6.55	6.23	6.13
MAX	5.91	6.30	6.40	6.40	6.37	7.04	6.60	---	7.60	6.91	6.33	6.27
MIN	5.75	5.75	6.27	6.31	6.31	6.42	6.43	---	6.99	6.28	6.13	5.95

434756089020500 GREEN LAKE AT DEEP HOLE NEAR GREEN LAKE, WI

LOCATION.--Lat 43°47'56", long 89°02'05", in NW 1/4 SE 1/4 sec.2, T.15 N., R.12 E., Green Lake County, Hydrologic Unit 04030201, about 5 miles southwest of the City of Green Lake.

PERIOD OF RECORD.--May to September 2004. Lake sampled by Wisconsin Department of Natural Resources prior to 2004.


REMARKS.--Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, MAY 18 TO JULY 23, 2004

(Milligrams per liter unless otherwise indicated)


Date	May-18	Jun-25	Jul-23
Lake stage (ft)	6.83	7.27	6.43
Secchi depth (m)	1.4	5.6	4.4
Depth of sample (m)	0.5	67.5	67
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	30.4	--	3.07
Water temperature ($^{\circ}\text{C}$)	11.5	5.3	23.5
Specific conductance ($\mu\text{S/cm}$)	485	505	489
pH	8.4	7.8	8.5
Dissolved oxygen (mg/L)	--	--	7.3
Phosphorus, total (as P)	0.037	0.049	0.017
Phosphorus, ortho, dissolved (as P)	0.002	--	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	<0.019	--	--
Nitrogen, ammonia, dissolved (as N)	<0.015	--	--
Nitrogen, amm. + org., total (as N)	0.68	--	--
Color (Pt-Co. scale)	10	--	--
Turbidity (NTU)	2.8	--	--
Hardness, as CaCO_3	210	--	--
Calcium, dissolved (Ca)	31.3	--	--
Magnesium, dissolved (Mg)	32.8	--	--
Sodium, dissolved (Na)	18.5	--	--
Potassium, dissolved (K)	3	--	--
Alkalinity, as CaCO_3	180	--	--
Sulfate, dissolved (SO_4)	30.5	--	--
Chloride, dissolved (Cl)	34.1	--	--
Silica, dissolved (SiO_2)	0.022	--	--
Solids, dissolved, at 180°C	272	--	--
Iron, dissolved (Fe) ($\mu\text{g/L}$)	<100	--	--
Lead, (Pb) ($\mu\text{g/L}$)	<1	--	--
Manganese, dissolved, (Mn) ($\mu\text{g/L}$)	<1	--	--

5-18-04




6-25-04


DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER





7-23-04



WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS




PH IN STANDARD UNITS







434756089020500 GREEN LAKE AT DEEP HOLE NEAR GREEN LAKE, WI--CONTINUED

WATER-QUALITY DATA, AUGUST 17 TO SEPTEMBER 16, 2004
 (Milligrams per liter unless otherwise indicated)

Date	Aug-17					Sep-16	
Lake stage (ft)		6.19				6.16	
Secchi depth (m)		3.0				3.5	
Depth of sample (m)	0.5	20	45	67	0.5	24	67
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	8.33	--	--	--	6.95	--	--
Water temperature ($^{\circ}\text{C}$)	20.7	7.8	6	5.5	20.5	6.9	5.6
Specific conductance ($\mu\text{S/cm}$)	487	499	503	510	478	496	510
pH	8.5	7.8	7.7	7.2	8.3	7.7	7.3
Dissolved oxygen (mg/L)	10.2	9.4	9.2	0.1	11.7	10.2	0.1
Phosphorus, total (as P)	0.019	0.019	0.06	0.165	0.012	0.015	0.211
Phosphorus, ortho, dissolved (as P)	<0.002	--	--	--	--	--	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	0.231	--	--	--	--	--	--
Nitrogen, ammonia, dissolved (as N)	0.017	--	--	--	--	--	--
Nitrogen, amm. + org., total (as N)	0.56	--	--	--	--	--	--
Nitrogen, total (as N)	0.79	--	--	--	--	--	--





Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Green Lake, Deep Hole, near Green Lake, Wisconsin.

434928088570000 GREEN LAKE AT EAST END NEAR GREEN LAKE, WI


LOCATION.--Lat 43°49'28", long 88°57'00", in SE 1/4 SE 1/4 sec.28, T.16 N., R.13 E., Green Lake County, Hydrologic Unit 04030201, about one mile southeast of the City of Green Lake.





PERIOD OF RECORD.--May to September 2004. Lake sampled by Wisconsin Department of Natural Resources prior to 2004.

REMARKS.--Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, MAY 18 TO SEPTEMBER 16, 2004
(Milligrams per liter unless otherwise indicated)

Date	May-18	Jun-25	Jul-23	Aug-17	Sep-16
Lake stage (ft)	6.83	7.27	6.43	6.19	6.16
Secchi depth (m)	1.0	4.7	2.7	2.7	2.7
Depth of sample (m)	0.5	34	0.5	34	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	54.5	--	1.4	--	--
Water temperature (°C)	12.1	5.8	19.0	6.4	21.0
Specific conductance ($\mu\text{S/cm}$)	487	500	485	506	477
pH	8.4	7.8	8.2	7.5	8.3
Dissolved oxygen (mg/L)	--	--	8.7	9.2	12.0
Phosphorus, total (as P)	0.047	0.055	0.044	0.067	0.016





Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Green Lake, East End, near Green Lake, Wisconsin.

05427235 LAKE KOSHKONONG NEAR NEWVILLE, WI

LOCATION.--Lat 42°51'27", long 88°56'27", in NW 1/4 NE 1/4 sec.34, T.5 N., R.13 E., Jefferson County, Hydrologic Unit 07090001, 80 ft east of Pottawatomi Trail Bridge at Bingham Point Estates, and 4.5 mi northeast of Newville.

DRAINAGE AREA.--2,560 mi², at lake outlet. Area of Lake Koshkonong, 16.3 mi².

PERIOD OF RECORD.--July 1987 to current year.

GAGE.--Water-stage recorder. Datum of gage is 770.00 ft above sea level.

REMARKS.--Lake level regulated by dam at Indianford. Gage-height telemeter at station.

EXTREMES FOR PERIOD OF RECORD.--Maximum recorded gage height, 12.23 ft, Apr. 25, 1993; minimum recorded, 5.10 ft, Dec. 28, 29, 1999.

EXTREMES FOR CURRENT YEAR.--Maximum recorded gage height, 11.49 ft, June 5; minimum daily gage height, 5.66 ft, Feb. 13.

GAGE HEIGHT, FEET
WATER YEAR OCTOBER 2003 TO SEPTEMBER 2004
DAILY MEAN VALUES

DAY	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP
1	6.12	5.90	6.42	5.85	5.80	6.01	8.16	6.70	11.12	10.60	6.84	6.31
2	6.09	5.94	6.37	5.80	5.81	6.09	8.16	6.63	11.30	10.46	6.75	6.29
3	6.09	6.02	6.34	5.81	5.85	6.20	8.17	6.55	11.42	10.34	6.66	6.27
4	6.10	6.22	6.30	5.81	5.90	6.34	8.08	6.47	11.48	10.30	6.76	6.27
5	6.08	6.34	6.26	5.75	5.91	6.65	8.01	6.39	11.49	10.17	6.78	6.26
6	6.08	6.44	6.21	5.75	5.88	6.96	7.96	6.31	11.46	10.04	6.77	6.26
7	6.08	6.57	6.14	5.76	5.84	7.29	7.91	6.21	11.40	9.95	6.73	6.24
8	6.08	6.63	6.10	5.77	5.80	7.59	7.86	6.18	11.33	9.83	6.70	6.21
9	6.08	6.70	6.07	5.77	5.75	7.82	7.78	6.25	11.23	9.71	6.69	6.24
10	6.08	6.79	6.20	5.77	5.71	7.99	7.71	6.31	11.16	9.61	6.66	6.26
11	6.08	6.90	6.23	5.78	5.68	8.10	7.61	6.45	11.17	9.49	6.59	6.30
12	6.11	7.04	6.24	5.78	5.67	8.17	7.51	6.54	11.17	9.38	6.47	6.33
13	6.09	7.06	6.27	5.78	5.66	8.21	7.42	6.69	11.20	9.28	6.36	6.33
14	6.16	6.97	6.31	5.78	5.68	8.27	7.31	6.85	11.28	9.17	6.27	6.30
15	6.14	6.92	6.35	5.76	5.70	8.25	7.19	6.96	11.32	9.05	6.27	6.27
16	6.13	6.88	6.33	5.74	5.71	8.24	7.10	7.05	11.34	8.94	6.26	6.30
17	6.12	6.82	6.29	5.74	5.72	8.22	7.03	7.15	11.44	8.82	6.29	6.26
18	6.12	6.83	6.23	5.72	5.73	8.20	6.92	7.28	11.45	8.68	6.29	6.23
19	6.12	6.76	6.17	5.70	5.74	8.16	6.88	7.35	11.43	8.55	6.27	6.20
20	6.13	6.68	6.08	5.70	5.77	8.16	6.75	7.42	11.39	8.42	6.23	6.19
21	6.15	6.59	5.99	5.69	5.80	8.07	6.84	7.54	11.37	8.31	6.22	6.21
22	6.13	6.49	5.94	5.69	5.82	7.99	6.86	8.03	11.36	8.22	6.20	6.23
23	6.13	6.48	5.94	5.69	5.86	7.94	6.88	8.57	11.29	8.09	6.21	6.24
24	6.13	6.51	5.92	5.69	5.90	7.91	6.88	9.11	11.26	7.94	6.25	6.26
25	6.18	6.43	5.91	5.68	5.91	7.87	6.91	9.58	11.19	7.81	6.29	6.27
26	6.19	6.44	5.90	5.67	5.89	7.94	6.92	9.96	11.12	7.67	6.30	6.27
27	6.17	6.43	5.91	5.68	5.90	7.94	6.85	10.25	11.03	7.51	6.35	6.27
28	6.12	6.46	5.95	5.71	5.91	7.97	6.78	10.45	10.96	7.36	6.37	6.29
29	6.05	6.40	5.97	5.74	5.94	8.06	6.77	10.60	10.84	7.21	6.37	6.25
30	5.99	6.42	5.95	5.76	---	8.14	6.74	10.75	10.72	7.07	6.36	6.25
31	5.96	---	5.91	5.78	---	8.15	--	10.93	---	6.95	6.33	---
MEAN	6.11	6.57	6.14	5.75	5.80	7.71	7.33	7.73	11.26	8.87	6.45	6.26
MAX	6.19	7.06	6.42	5.85	5.94	8.27	8.17	10.93	11.49	10.60	6.84	6.33
MIN	5.96	5.90	5.90	5.67	5.66	6.01	6.74	6.18	10.72	6.95	6.20	6.19

432255088134700 LITTLE CEDAR LAKE, NORTH SITE, NEAR WEST BEND, WI


LOCATION.--Lat 43°22'55", long 88°13'47", in NW 1/4 NE 1/4 sec.33, T.11 N., R.19 E., Washington County, Hydrologic Unit 04040003, 2.6 mi southwest of West Bend.


PERIOD OF RECORD.--February 1997 to August 1999, February 2003 to current year.

REMARKS.--Lake sampled at center of northern basin at deep hole. Lake ice-covered during February sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, FEBRUARY 24 TO AUGUST 18, 2004
(Milligrams per liter unless otherwise indicated)

Date	<u>Feb-24</u>	<u>Apr-22</u>	<u>Jun-9</u>	<u>Jul-14</u>	<u>Aug-18</u>
Lake stage (ft)	--	--	4.86	4.98	--
Secchi depth (m)	--	1.0	2.2	2.0	1.5
Depth of sample (m)	0.5	6.5	0.5	7.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	--	28.0	9.0	--
Water temperature ($^{\circ}\text{C}$)	0.6	4.0	12.0	9.8	24.2
Specific conductance ($\mu\text{S/cm}$)	589	620	538	545	531
pH	7.6	7.2	8.5	8.1	7.4
Dissolved oxygen (mg/L)	12.3	4.3	10.9	9.0	11.4
Phosphorus, total (as P)	0.015	0.025	0.041	0.035	0.028





432249088134500 LITTLE CEDAR LAKE, SOUTH SITE, NEAR WEST BEND, WI


LOCATION.--Lat 43°22'49", long 88°13'45", in NW 1/4 SE 1/4 sec.33, T.11 N., R.19 E., Washington County, Hydrologic Unit 04040003, 2.8 mi southwest of West Bend.


PERIOD OF RECORD.--February 1997 to current year.

REMARKS.--Lake sampled in southern basin at deep hole. Lake ice-covered during February sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, FEBRUARY 24 TO AUGUST 18, 2004
(Milligrams per liter unless otherwise indicated)

Date	Feb-24	Apr-22	Jun-9	Jul-14	Aug-18
Lake stage (ft)	--	--	4.86	4.98	--
Secchi depth (m)	--	1.0	2.1	2.8	1.8
Depth of sample (m)	0.5	16.5	0.5	0.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	--	33.4	--	8.4
Water temperature (°C)	0.8	4.2	11.5	6.6	21.3
Specific conductance ($\mu\text{S/cm}$)	560	623	532	547	508
pH	8.0	7.1	8.5	7.7	7.5
Dissolved oxygen (mg/L)	15.0	0.6	11.6	4.7	8.8
Phosphorus, total (as P)	0.031	0.113	0.038	0.059	0.040
Phosphorus, ortho, dissolved (as P)	--	--	0.003	--	<0.002
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	--	--	<0.019	--	<0.019
Nitrogen, ammonia, dissolved (as N)	--	--	<0.015	--	<0.015
Nitrogen, amm. + diss., total (as N)	--	--	--	--	0.64
Nitrogen, amm. + org., total (as N)	--	--	0.83	--	--
Color (Pt-Co. scale)	--	--	15	--	--
Turbidity (NTU)	--	--	4.7	--	--
Hardness, as CaCO_3	--	--	220	--	--
Calcium, dissolved (Ca)	--	--	34.6	--	--
Magnesium, dissolved (Mg)	--	--	32.2	--	--
Sodium, dissolved (Na)	--	--	21	--	--
Potassium, dissolved (K)	--	--	2	--	--
Alkalinity, as CaCO_3	--	--	185	--	--
Sulfate, dissolved (SO_4)	--	--	18	--	--
Chloride, dissolved (Cl)	--	--	45.7	--	--
Silica, dissolved (SiO_2)	--	--	2.97	--	--
Solids, dissolved, at 180°C	--	--	300	--	--
Iron, dissolved (Fe) ($\mu\text{g/L}$)	--	--	<100	--	--
Manganese, dissolved, (Mn) ($\mu\text{g/L}$)	--	--	<1	--	--





05428000 LAKE MENDOTA AT MADISON, WI

LOCATION.--Lat 43°05'42", long 89°22'12", in SE 1/4 sec.12, T.7 N., R.9 E., Dane County, Hydrologic Unit 07090001, in county boat house at dam at outlet, in Madison.

DRAINAGE AREA.--233 mi². Area of Lake Mendota, 15.2 mi².

PERIOD OF RECORD.--January 1916 to current year (incomplete).

REVISED RECORDS.--WDR WI-73-1: Drainage area.

GAGE.--Water-stage recorder. Datum of gage is 840.00 ft above sea level, or 5.60 ft below City of Madison datum. Prior to Oct. 1, 1979, at datum 7.82 ft higher; prior to Nov. 15, 1971, nonrecording gage at same site.

REMARKS.--Lake level regulated by concrete dam with two 12-foot gates and 20-foot lock at outlet. Gage-height telemeter at station.

EXTREMES FOR PERIOD OF RECORD.--Maximum gage height observed, 12.75 ft, June 5, 2000; minimum observed, 8.02 ft, Feb. 24 to Mar. 10, 1920, current datum.

EXTREMES FOR CURRENT YEAR.--Maximum recorded gage height, 11.61 ft, May 26; minimum recorded, 8.87 ft, Feb. 18, 19.

GAGE HEIGHT, FEET
WATER YEAR OCTOBER 2003 TO SEPTEMBER 2004
DAILY MEAN VALUES

DAY	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP
1	9.72	9.26	9.56	9.39	9.01	9.16	9.99	9.74	11.57	10.81	10.26	10.00
2	9.68	9.33	9.49	9.39	9.01	9.22	9.96	9.73	11.53	10.79	10.30	10.01
3	9.67	9.43	9.44	9.39	9.01	9.26	9.96	9.72	11.48	10.78	10.34	10.03
4	9.66	9.84	9.43	9.36	9.00	9.30	9.89	9.71	11.41	11.03	10.49	10.02
5	9.64	9.98	9.44	9.35	9.00	9.48	9.85	9.70	11.34	11.02	10.49	10.00
6	9.63	10.04	9.42	9.30	9.02	9.60	9.84	9.70	11.28	10.96	10.49	9.98
7	9.63	10.07	9.39	9.29	9.01	9.66	9.82	9.69	11.21	10.95	10.47	9.93
8	9.64	10.06	9.39	9.29	8.99	9.69	9.80	9.69	11.16	10.92	10.43	9.88
9	9.65	10.03	9.39	9.27	8.98	9.70	9.75	9.76	11.11	10.85	10.43	9.84
10	9.64	9.99	9.56	9.26	8.97	9.70	9.72	9.81	11.08	10.81	10.38	9.81
11	9.65	9.98	9.59	9.26	8.96	9.70	9.67	9.83	11.17	10.75	10.31	9.79
12	9.66	10.00	9.57	9.24	8.95	9.70	9.64	9.85	11.22	10.72	10.24	9.78
13	9.64	9.93	9.56	9.23	8.94	9.70	9.63	9.89	11.25	10.68	10.21	9.76
14	9.67	9.87	9.56	9.22	8.93	9.71	9.62	9.94	11.25	10.61	10.18	9.75
15	9.63	9.84	9.55	9.21	8.91	9.72	9.61	9.94	11.22	10.54	10.15	9.74
16	9.59	9.81	9.59	9.20	8.90	9.71	9.61	9.90	11.19	10.50	10.10	9.76
17	9.55	9.77	9.56	9.20	8.89	9.72	9.66	9.94	11.20	10.48	10.09	9.74
18	9.52	9.81	9.55	9.18	8.87	9.73	9.68	10.02	11.17	10.42	10.09	9.72
19	9.50	9.82	9.54	9.17	8.87	9.73	9.71	10.02	11.08	10.38	10.08	9.69
20	9.48	9.76	9.51	9.16	8.89	9.75	9.70	10.05	11.01	10.34	10.05	9.67
21	9.46	9.71	9.50	9.14	8.90	9.73	9.77	10.14	10.96	10.32	10.02	9.67
22	9.41	9.66	9.50	9.11	8.90	9.72	9.77	10.67	10.92	10.29	10.00	9.66
23	9.39	9.81	9.49	9.11	8.91	9.72	9.76	11.16	10.88	10.22	10.00	9.64
24	9.38	9.86	9.46	9.11	8.92	9.75	9.76	11.46	10.91	10.17	9.98	9.62
25	9.41	9.81	9.44	9.10	8.93	9.78	9.80	11.58	10.90	10.15	10.01	9.59
26	9.38	9.79	9.43	9.09	8.96	9.93	9.81	11.61	10.88	10.12	9.97	9.58
27	9.35	9.75	9.42	9.08	9.00	9.96	9.78	11.60	10.86	10.10	10.00	9.58
28	9.33	9.72	9.44	9.07	9.04	10.01	9.77	11.56	10.87	10.08	9.99	9.55
29	9.31	9.64	9.44	9.05	9.09	10.05	9.77	11.52	10.85	10.05	9.97	9.53
30	9.29	9.61	9.42	9.03	--	10.06	9.76	11.54	10.82	10.17	9.95	9.53
31	9.30	---	9.41	9.02	--	10.02	--	11.60	---	10.24	9.97	---
MEAN	9.53	9.80	9.49	9.20	8.96	9.70	9.76	10.36	11.13	10.52	10.18	9.76
MAX	9.72	10.07	9.59	9.39	9.09	10.06	9.99	11.61	11.57	11.03	10.49	10.03
MIN	9.29	9.26	9.39	9.02	8.87	9.16	9.61	9.69	10.82	10.05	9.95	9.53

424621088335500 MIDDLE LAKE AT LAUDERDALE, WI

LOCATION.--Lat 42°46'21", long 88°33'55", in SE 1/4 SE 1/4 sec.26, T.4 N., R.16 E., Walworth County, Hydrologic Unit 07120006, at Lauderdale.
 PERIOD OF RECORD.--November 1993 to November 1994, February 1999 to current year.

REMARKS.--Lake sampled near east end of lake at a depth of about 52 ft. Lake ice-covered during February sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, FEBRUARY 18 TO AUGUST 16, 2004
 (Milligrams per liter unless otherwise indicated)

Date	Feb-18	Apr-8	Jun-7	Jul-12	Aug-16
Secchi depth (m)	--	--	4.3	2.7	2.2
Depth of sample (m)	0.5	13.5	0.5	14	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	--	2.0	--	4.8
Water temperature ($^{\circ}\text{C}$)	0.6	4.5	10.1	4.9	21.6
Specific conductance ($\mu\text{S/cm}$)	540	662	521	619	491
pH	8.0	7.6	8.3	7.6	8.0
Dissolved oxygen (mg/L)	15.1	8.4	11.9	7.6	9.3
Phosphorus, total (as P)	0.011	0.012	0.015	0.021	0.047
Phosphorus, ortho, dissolved (as P)	--	--	0.001	--	<0.002
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	--	--	0.868	--	0.309
Nitrogen, ammonia, dissolved (as N)	--	--	0.223	--	0.08
Nitrogen, amm. + diss., total (as N)	--	--	--	--	0.65
Nitrogen, amm. + org., total (as N)	--	--	--	--	--
Nitrogen, total (as N)	--	--	0.89	--	--
Color (Pt-Co. scale)	--	--	5	--	--
Turbidity (NTU)	--	--	1.2	--	--
Hardness, as CaCO_3	--	--	250	--	--
Calcium, dissolved (Ca)	--	--	44.2	--	--
Magnesium, dissolved (Mg)	--	--	34.1	--	--
Sodium, dissolved (Na)	--	--	8.8	--	--
Potassium, dissolved (K)	--	--	2	--	--
Alkalinity, as CaCO_3	--	--	202	--	--
Sulfate, dissolved (SO_4)	--	--	36.1	--	--
Chloride, dissolved (Cl)	--	--	24.3	--	--
Silica, dissolved (SiO_2)	--	--	2.85	--	--
Solids, dissolved, at 180°C	--	--	292	--	--
Iron, dissolved (Fe) ($\mu\text{g/L}$)	--	--	<100	--	--
Manganese, dissolved, (Mn) ($\mu\text{g/L}$)	--	--	<1	--	--

2-18-04


4-8-04

6-7-04

7-12-04


8-16-04


DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER



WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS

PH IN STANDARD UNITS





430251088284700 MIDDLE GENESEE LAKE NEAR OCONOMOWOC, WI

LOCATION.--Lat 43°02'51", long 88°28'47", in SW 1/4 SW 1/4 SW 1/4 sec.22, T. 7 N., R.17 E., Waukesha County, Hydrologic Unit 07090001, at the southwest side of the lake about 2 miles south of Oconomowoc.

DRAINAGE AREA.--Unknown. Area of Middle Genesee Lake is 0.17 mi².

PERIOD OF RECORD.--April 1996 to current year.

GAGE.--Staff gage. Local observer, Tom Schubring provided most readings of gage.


EXTREMES FOR THE PERIOD OF RECORD.--Maximum observed gage height, 867.18 ft, June 13, 2001; minimum observed, 863.92 ft, Oct. 22, 31 and Nov. 1, 2003.

EXTREMES FOR CURRENT YEAR.--Maximum observed gage height, 867.14 ft, June 22; minimum observed, 863.92 ft, Oct. 22, 31 and Nov. 1.

GAGE HEIGHT, FEET, WATER YEAR OCTOBER 2003 TO OCTOBER 2004

DAILY MEAN VALUES

Date	Gage height (feet)	Date	Gage height (feet)	Date	Gage height (feet)	Date	Gage height (feet)
October 1	864.08	19	864.34	June 24	867.12	August 14	866.90
3	864.06	29	864.34	27	867.09	20	866.84
8	864.02	December 2	864.32	July 4	867.06	31	866.86
11	864.00	8	864.32	9	867.02	September 6	866.78
18	863.98	April 14	864.86	12	867.07	12	866.68
22	863.92	22	865.20	15	867.02	18	866.58
26	863.98	May 2	865.22	18	867.02	26	866.44
29	863.94	9	865.26	July 26	866.90	30	866.36
31	863.92	10	865.40	29	866.88	October 4	866.26
November 1	863.92	26	866.52	31	866.84	15	866.08
4	864.28	29	866.62	August 1	866.82	24	865.94
5	864.30	31	866.76	3	866.84	28	865.86
14	864.26	June 8	867.00	6	867.04	31	865.82
17	864.26	22	867.14	8	867.00		



430309088284800 MIDDLE GENESEE LAKE NEAR OCONOMOWOC, WI

LOCATION.--Lat 43°03'09", long 88°28'48", in NW 1/4 SW 1/4 sec.22, T.7 N., R.17 E., Waukesha County, Hydrologic Unit 07090001, 1.8 mi south of Oconomowoc.

PERIOD OF RECORD.--February 1996 to current year.

REMARKS.--Lake sampled near center at the deep hole. Lake ice-covered during February sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, FEBRUARY 19 TO AUGUST 3, 2004
(Milligrams per liter unless otherwise indicated)

Date	<u>Feb-19</u>	<u>Apr-14</u>	<u>Jun-8</u>	<u>Jul-15</u>	<u>Aug-3</u>
Lake stage (ft)	--	864.86	867.00	876.02	866.84
Secchi depth (m)	--	8.4	3.4	3.5	3.2
Depth of sample (m)	0.5	11.5	0.5	12	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	0.8	--	3.7	--
Water temperature (°C)	2.7	5.4	9.4	8.3	22.5
Specific conductance ($\mu\text{S/cm}$)	477	555	421	420	418
pH	8.0	7.2	8.2	8.2	455
Dissolved oxygen (mg/L)	13.4	0.9	11.8	11.3	9.9
Phosphorus, total (as P)	0.009	0.017	0.014	0.015	0.023
Phosphorus, ortho, dissolved (as P)	--	<0.002	--	--	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	--	--	0.042	--	--
Nitrogen, ammonia, dissolved (as N)	--	--	0.197	--	--
Nitrogen, amm. + org., total (as N)	--	--	0.89	--	--
Nitrogen, total (as N)	--	--	0.93	--	--
Color (Pt-Co. scale)	--	--	5	--	--
Turbidity (NTU)	--	--	<1	--	--
Hardness, as CaCO_3	--	--	190	--	--
Calcium, dissolved (Ca)	--	--	30.8	--	--
Magnesium, dissolved (Mg)	--	--	27.6	--	--
Sodium, dissolved (Na)	--	--	13.1	--	--
Potassium, dissolved (K)	--	--	1	--	--
Alkalinity, as CaCO_3	--	--	160	--	--
Sulfate, dissolved (SO_4)	--	--	14.8	--	--
Chloride, dissolved (Cl)	--	--	28.9	--	--
Silica, dissolved (SiO_2)	--	--	1.14	--	--
Solids, dissolved, at 180°C	--	--	236	--	--
Iron, dissolved (Fe) ($\mu\text{g/L}$)	--	--	<100	--	--
Manganese, dissolved (Mn) ($\mu\text{g/L}$)	--	--	<1	--	--

2-19-04


4-14-04

6-8-04

7-15-04


8-3-04

DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER




WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS

PH IN STANDARD UNITS



SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS



Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Middle Genesee Lake, near Oconomowoc, Wisconsin.

(Circles on the first two plots indicate laboratory detection limit for selected analyses.
Actual concentrations for these particular analyses are less than the plotted circles.)

05429000 LAKE MONONA AT MADISON, WI

LOCATION.--Lat 43°03'48", long 89°23'49", in SE 1/4 SW 1/4 sec.23, T.7 N., R.9 E., Dane County, Hydrologic Unit 07090001, in Brittingham Park, in Madison.

DRAINAGE AREA.--279 mi². Area of Lake Monona, 5.3 mi².

PERIOD OF RECORD.--September 1915 to current year (fragmentary) in reports of the Geological Survey. For 1856 to March 1917 in reports of Wisconsin Railroad Commission, volume 19.

REVISED RECORDS.--WSP 1338: Lake area. WDR WI-73-1: Drainage area.

GAGE.--Water-stage recorder. Datum of gage is 840.00 ft above sea level, or 5.60 ft below City of Madison datum. Prior to Oct. 1, 1979, datum 3.61 ft higher; prior to Nov. 15, 1971, nonrecording gage at same site at the higher datum.

REMARKS.--Lake level regulated by concrete dam with four 12-foot stop-log sections and 12-foot lock at outlet of Lake Waubesa. Gage-height telemeter at station.

EXTREMES FOR PERIOD OF RECORD.--Maximum gage height observed, 7.48 ft, June 14, 15, 2000; minimum observed, 3.22 ft, Jan. 20, 1965, current datum.

EXTREMES FOR CURRENT YEAR.--Maximum recorded gage height, 6.87 ft, June 11; minimum recorded, 4.10, Feb. 17, 18, 19.

**GAGE HEIGHT, FEET
WATER YEAR OCTOBER 2003 TO SEPTEMBER 2004
DAILY MEAN VALUES**

DAY	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP
1	4.84	4.52	5.39	4.48	4.15	4.19	4.91	4.68	6.75	6.31	6.07	5.82
2	4.83	4.70	5.38	4.47	4.15	4.19	4.94	4.68	6.75	6.34	6.02	5.83
3	4.81	4.91	5.37	4.45	4.16	4.18	4.93	4.68	6.73	6.40	6.01	5.83
4	4.79	5.36	5.28	4.43	4.16	4.18	4.94	4.68	6.73	6.61	6.17	5.83
5	4.79	5.47	5.21	4.43	4.15	4.38	4.97	4.68	6.73	6.62	6.15	5.82
6	4.78	5.40	5.14	4.43	4.17	4.45	4.99	4.69	6.73	6.58	6.11	5.82
7	4.77	5.30	5.10	4.42	4.17	4.45	5.01	4.72	6.73	6.54	6.08	5.81
8	4.76	5.23	5.06	4.40	4.17	4.42	5.00	4.71	6.73	6.50	6.07	5.80
9	4.75	5.22	5.02	4.39	4.16	4.41	5.01	4.82	6.74	6.52	6.04	5.77
10	4.75	5.23	5.15	4.37	4.15	4.41	5.01	4.89	6.76	6.53	6.01	5.76
11	4.75	5.22	5.16	4.36	4.15	4.39	5.01	4.99	6.87	6.51	5.97	5.74
12	4.74	5.17	5.14	4.34	4.13	4.36	5.01	5.04	6.83	6.50	5.94	5.72
13	4.74	5.12	5.12	4.31	4.13	4.36	4.96	5.10	6.76	6.46	5.93	5.71
14	4.81	5.12	5.08	4.29	4.12	4.35	4.94	5.13	6.70	6.43	5.92	5.67
15	4.80	5.12	5.04	4.28	4.11	4.34	4.92	5.15	6.65	6.42	5.91	5.66
16	4.79	5.12	4.98	4.26	4.11	4.33	4.88	5.15	6.63	6.47	5.89	5.66
17	4.77	5.12	4.93	4.27	4.10	4.34	4.91	5.15	6.72	6.58	5.90	5.62
18	4.76	5.16	4.89	4.25	4.10	4.34	4.89	5.26	6.72	6.56	5.89	5.57
19	4.76	5.13	4.84	4.23	4.10	4.34	4.87	5.24	6.71	6.52	5.91	5.54
20	4.74	5.16	4.79	4.22	4.14	4.31	4.87	5.20	6.68	6.49	5.88	5.50
21	4.71	5.20	4.75	4.22	4.17	4.29	4.88	5.29	6.68	6.48	5.84	5.44
22	4.69	5.23	4.72	4.20	4.19	4.30	4.87	5.86	6.66	6.47	5.82	5.39
23	4.65	5.39	4.69	4.20	4.21	4.32	4.84	6.40	6.63	6.45	5.81	5.34
24	4.65	5.45	4.65	4.20	4.23	4.35	4.82	6.56	6.59	6.39	5.78	5.28
25	4.69	5.47	4.62	4.19	4.23	4.38	4.81	6.61	6.55	6.33	5.79	5.24
26	4.65	5.48	4.60	4.19	4.22	4.59	4.76	6.60	6.49	6.27	5.79	5.20
27	4.62	5.48	4.58	4.19	4.20	4.67	4.73	6.57	6.44	6.21	5.81	5.16
28	4.59	5.44	4.58	4.18	4.19	4.71	4.72	6.56	6.40	6.11	5.83	5.11
29	4.57	5.44	4.55	4.17	4.18	4.76	4.71	6.60	6.32	6.04	5.83	5.06
30	4.56	5.42	4.53	4.16	---	4.81	4.70	6.67	6.29	6.10	5.82	5.02
31	4.52	---	4.50	4.16	---	4.87	---	6.72	---	6.12	5.81	---
MEAN	4.72	5.23	4.93	4.29	4.16	4.41	4.89	5.45	6.66	6.41	5.93	5.56
MAX	4.84	5.48	5.39	4.48	4.23	4.87	5.01	6.72	6.87	6.62	6.17	5.83
MIN	4.52	4.52	4.50	4.16	4.10	4.18	4.70	4.68	6.29	6.04	5.78	5.02

425109088075000 MUSKEGO (BIG MUSKEGO) LAKE NEAR WIND LAKE, WI

LOCATION.--Lat 42°51'09", long 88°07'50", in SE 1/4 NE 1/4 sec.33, T.5 N., R.20 E., Waukesha County, Hydrologic Unit 07120006, on left bank 8 ft upstream of dam outlet of Muskego Lake, 700 ft north of Muskego Dam Drive, 2 mi northeast of Wind Lake.

DRAINAGE AREA.--33.9 mi² (revised).

PERIOD OF RECORD.--October 1987 to September 1989, January 1991 to current year. Prior to October 1993, published as Muskego Lake Outlet near Wind Lake, WI. October 1993 to September 2000, published as "Big Muskego Lake".

GAGE.--Water-stage recorder. Datum of gage is 760.00 ft above sea level. October to December 1987 and January 1991 to September 1995, nonrecording gage at the same datum. December 1987 through September 1989, data collected using water-stage recorder at the same datum.

REMARKS.--Lake levels regulated by concrete dam with one 5-ft lift gate.

EXTREMES FOR PERIOD OF RECORD.--Maximum observed gage height, 12.60 ft, Oct. 7, 1991 and Aug. 8, 1994; minimum instantaneous, less than 8.72 ft, July 12, 1996 to Feb. 18, 1997, due to drawdown of lake.

EXTREMES FOR CURRENT YEAR.--Maximum observed gage-height, 12.25 ft, May 25; minimum observed, 10.58 ft, Oct. 20.

GAGE HEIGHT, FEET
WATER YEAR OCTOBER 2003 TO SEPTEMBER 2004
DAILY MEAN VALUES

DAY	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP
1	10.71	10.99	11.58	11.70	11.69	11.73	11.82	e11.44	11.96	11.47	11.00	11.09
2	10.67	11.08	11.60	11.70	11.69	11.77	11.80	e11.42	11.97	11.46	11.00	11.08
3	10.61	11.25	11.60	11.70	11.70	11.78	11.84	e11.42	11.90	11.44	11.03	11.09
4	10.69	11.28	11.61	11.70	11.70	11.81	11.76	e11.38	11.82	11.57	11.11	11.08
5	10.69	11.33	11.61	11.70	11.69	11.94	11.72	e11.32	11.74	11.64	11.11	11.07
6	10.67	11.34	11.61	11.70	11.71	11.97	11.71	e11.28	11.67	11.61	11.07	11.02
7	10.64	11.36	11.61	11.69	11.70	11.95	11.70	11.23	11.63	11.69	11.06	11.08
8	10.64	11.40	11.62	11.69	11.70	11.93	11.63	11.10	11.62	11.64	11.04	11.10
9	10.66	11.40	11.64	11.69	11.69	11.89	11.55	11.15	11.62	11.54	11.02	11.06
10	10.66	11.40	11.80	11.69	11.68	11.82	11.50	11.16	11.62	11.50	11.02	11.03
11	10.62	11.40	11.77	11.69	11.68	11.78	11.44	11.21	11.65	11.43	11.02	11.01
12	10.63	11.32	11.77	11.68	11.67	11.77	11.41	11.15	11.65	11.40	11.04	11.01
13	10.61	11.41	11.77	11.68	11.67	11.73	11.37	11.30	11.62	11.31	11.03	11.00
14	10.71	11.38	11.77	11.68	11.66	11.64	11.27	11.49	11.62	11.25	11.02	10.97
15	10.65	11.41	11.76	11.68	11.65	11.71	11.19	11.60	11.64	11.20	11.00	10.89
16	10.67	11.43	11.75	11.68	11.65	11.65	11.16	11.60	11.61	11.25	10.97	10.99
17	10.63	11.44	11.75	11.69	11.64	11.57	11.25	11.56	11.63	11.32	10.99	11.00
18	10.61	11.53	11.74	11.69	11.64	11.56	11.01	11.70	11.62	11.26	10.95	10.97
19	10.65	11.55	11.73	11.68	11.64	11.51	11.02	11.69	11.61	11.22	10.99	10.94
20	10.58	11.53	11.72	11.68	11.65	11.51	11.17	11.68	11.53	11.21	10.98	10.93
21	10.67	11.64	11.71	11.68	11.66	11.52	11.18	11.73	11.55	11.20	10.97	10.91
22	10.69	11.63	11.71	11.68	11.65	11.41	11.34	11.85	11.59	11.21	10.89	10.92
23	10.70	11.53	11.70	11.69	11.66	11.42	11.28	12.08	11.56	11.20	10.97	10.90
24	10.73	11.49	11.70	11.70	11.67	11.43	11.51	12.23	11.57	11.15	10.94	10.85
25	10.82	11.58	11.69	11.69	11.67	11.43	11.49	12.25	11.54	11.14	11.02	10.89
26	10.85	11.60	11.69	11.70	11.67	11.59	e11.46	12.21	11.52	11.09	11.01	10.88
27	10.87	11.61	11.69	11.70	11.68	11.65	e11.44	12.15	11.49	11.07	11.03	10.87
28	10.83	11.61	11.70	11.70	11.68	11.63	e11.42	12.09	11.49	11.05	11.11	11.00
29	10.90	11.60	11.71	11.70	11.70	11.76	e11.44	12.01	11.43	11.03	11.11	10.86
30	10.94	11.59	11.70	11.70	---	11.89	e11.44	11.99	11.44	11.03	11.09	10.83
31	10.94	---	11.70	11.70	---	11.86	---	11.99	---	11.03	11.10	---
MEAN	10.71	11.44	11.69	11.69	11.67	11.70	11.44	11.63	11.63	11.31	11.02	10.98
MAX	10.94	11.64	11.80	11.70	11.71	11.97	11.84	12.25	11.97	11.69	11.11	11.10
MIN	10.58	10.99	11.58	11.68	11.64	11.41	11.01	11.10	11.43	11.03	10.89	10.83

e Estimated

430347088240800 NAGAWICKA LAKE AT DELAFIELD, WI

LOCATION.--Lat 43°03'47", long 88°24'08", in SW 1/4 SW 1/4 sec.17, T.7 N., R.18 E., Waukesha County, Hydrologic Unit 07090001, on dike of Nagawicka Lake dam about 120 ft west of gates in Delafield.

DRAINAGE AREA.--44.9 mi². Area of Nagawicka Lake, 917 acres.

PERIOD OF RECORD.--October 2002 to current year.

GAGE.--Water-stage recorder.

REMARKS.--Gage established Oct. 29, 2002. Lake levels controlled by City of Delafield.

EXTREMES FOR CURRENT YEAR.--Maximum gage height, 8.73 ft, May 23, affected by wind; minimum gage height, 7.60 ft, Dec. 1.

**GAGE HEIGHT, FEET
WATER YEAR OCTOBER 2003 TO SEPTEMBER 2004
DAILY MEAN VALUES**

DAY	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP
1	8.28	8.35	7.64	7.92	7.80	7.84	8.27	8.38	8.41	8.42	8.34	8.37
2	8.27	8.44	7.67	7.92	7.79	7.89	8.24	8.38	8.47	8.42	8.34	8.37
3	8.28	8.55	7.70	7.92	7.80	7.93	8.23	8.36	8.49	8.40	8.40	8.37
4	8.30	8.61	7.72	7.93	7.79	7.97	8.20	8.34	8.48	8.40	8.56	8.36
5	8.32	8.51	7.74	7.92	7.79	8.12	8.18	8.35	8.45	8.41	8.42	8.35
6	8.32	8.35	7.74	7.90	7.80	8.14	8.19	8.35	8.41	8.38	8.36	8.34
7	8.32	8.23	7.75	7.89	7.80	8.15	8.21	8.38	8.38	8.40	8.36	8.34
8	8.33	8.24	7.76	7.89	7.79	8.14	8.21	8.35	8.41	8.41	8.36	8.33
9	8.34	8.24	7.80	7.88	7.79	8.12	8.21	8.39	8.43	8.44	8.37	8.31
10	8.34	8.25	7.93	7.88	7.78	8.11	8.22	8.40	8.46	8.46	8.37	8.29
11	8.34	8.22	7.96	7.87	7.77	8.11	8.23	8.52	8.51	8.47	8.36	8.28
12	8.36	8.18	7.97	7.87	7.77	8.11	8.23	8.54	8.50	8.55	8.37	8.27
13	8.35	8.15	8.00	7.86	7.76	8.10	8.22	8.56	8.49	8.51	8.36	8.26
14	8.39	8.15	8.02	7.86	7.76	8.10	8.21	8.52	8.48	8.47	8.35	8.25
15	8.36	8.16	8.01	7.85	7.75	8.11	8.21	8.48	8.49	8.43	8.35	8.23
16	8.33	8.16	7.90	7.85	7.74	8.10	8.21	8.41	8.48	8.42	8.34	8.23
17	8.30	8.16	7.90	7.87	7.74	8.09	8.29	8.34	8.55	8.44	8.36	8.22
18	8.27	8.22	7.90	7.86	7.73	8.09	8.31	8.42	8.49	8.42	8.36	8.21
19	8.26	8.24	7.90	7.85	7.73	8.08	8.34	8.40	8.46	8.40	8.38	8.19
20	8.23	8.25	7.88	7.84	7.74	8.07	8.40	8.34	8.42	8.38	8.38	8.17
21	8.24	8.27	7.88	7.83	7.76	8.07	8.51	8.42	8.45	8.38	8.39	8.15
22	8.25	8.28	7.89	7.82	7.76	8.05	8.45	8.61	8.48	8.40	8.38	8.15
23	8.25	8.25	7.89	7.82	7.77	8.05	8.36	8.67	8.46	8.39	8.39	8.17
24	8.27	8.15	7.90	7.83	7.78	8.07	8.40	8.69	8.47	8.37	8.36	8.17
25	8.33	8.08	7.89	7.82	7.78	8.10	8.43	8.69	8.46	8.35	8.37	8.20
26	8.33	8.01	7.89	7.83	7.78	8.22	8.43	8.67	8.46	8.34	8.38	8.21
27	8.33	7.94	7.89	7.83	7.79	8.23	8.39	8.63	8.45	8.34	8.46	8.23
28	8.33	7.85	7.91	7.83	7.80	8.24	8.33	8.57	8.45	8.33	8.48	8.25
29	8.33	7.76	7.92	7.82	7.81	8.28	8.33	8.49	8.43	8.33	8.43	8.23
30	8.35	7.68	7.92	7.81	---	8.30	8.37	8.45	8.42	8.34	8.37	8.24
31	8.35	---	7.91	7.80	---	8.28	---	8.42	---	8.34	8.37	---
MEAN	8.31	8.20	7.86	7.86	7.77	8.11	8.29	8.47	8.46	8.40	8.38	8.26
MAX	8.39	8.61	8.02	7.93	7.81	8.30	8.51	8.69	8.55	8.55	8.56	8.37
MIN	8.23	7.68	7.64	7.80	7.73	7.84	8.18	8.34	8.38	8.33	8.34	8.15

430417088230300 NAGAWICKA LAKE, AT DEEP HOLE, AT DELAFIELD, WI

LOCATION.--Lat 43°04'17", long 88°23'03", in SE 1/4 NE 1/4 sec.17, T.7 N., R.18 E., Waukesha County, Hydrologic Unit 07090001, at Delafield.

PERIOD OF RECORD.--February 2003 to current year.


REMARKS.--Lake sampled near center at the deep hole. Lake ice-covered during February sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, FEBRUARY 19 TO JUNE 7, 2004

(Milligrams per liter unless otherwise indicated)


Date	Feb-19	Apr-14	Jun-7
Lake stage (ft)	7.73	8.21	8.38
Secchi depth (m)	--	5.0	5.5
Depth of sample (m)	0.5	26.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	--	1.7
Water temperature ($^{\circ}\text{C}$)	0.3	1.8	6.0
Specific conductance ($\mu\text{S/cm}$)	804	877	731
pH	8.0	7.5	8.1
Dissolved oxygen (mg/L)	14.2	8.1	12.3
Phosphorus, total (as P)	0.008	0.016	0.012
Phosphorus, ortho, dissolved (as P)	--	<0.002	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	--	0.942	--
Nitrogen, ammonia, dissolved (as N)	--	--	--
Nitrogen, amm. + org., total (as N)	--	0.098	--
Nitrogen, total (as N)	--	--	0.58
Color (Pt-Co. scale)	--	--	1.5
Turbidity (NTU)	--	--	<1.0
Hardness, as CaCO_3	--	--	300
Calcium, dissolved (Ca)	--	--	56.7
Magnesium, dissolved (Mg)	--	--	38.3
Sodium, dissolved (Na)	--	--	33.4
Potassium, dissolved (K)	--	--	2
Chloride, dissolved (Cl)	--	--	74.1
Solids, dissolved, at 180°C	--	--	414
Iron, dissolved (Fe) ($\mu\text{g/L}$)	--	--	<100
Manganese, dissolved, (Mn) ($\mu\text{g/L}$)	--	--	<1

2-19-04




4-14-04


DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER





6-7-04



WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS



PH IN STANDARD UNITS




430417088230300 NAGAWICKA LAKE, AT DEEP HOLE, AT DELAFIELD, WI--CONTINUED

WATER-QUALITY DATA, JULY 14 TO SEPTEMBER 23, 2004
(Milligrams per liter unless otherwise indicated)


Date	Jul-14	Aug-17				Sep-23				
Lake stage (ft)	8.47		8.36				8.17			
Secchi depth (m)	4.2		5.1				2.7			
Depth of sample (m)	0.5	27.5	0.5	9	26.5	0.5	8	17	20	23
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	5.9	--	3.3	--	--	4.8	--	--	--	--
Water temperature ($^{\circ}\text{C}$)	23.9	7.8	20.7	15.8	8.0	21.0	20.0	8.8	8.3	8.3
Specific conductance ($\mu\text{S/cm}$)	702	723	705	711	723	699	704	740	742	744
pH	8.2	7.6	8.2	7.6	7.6	8.5	8.4	9.1	9.0	8.7
Dissolved oxygen (mg/L)	11.9	0.3	11.2	1.9	0.3	9.0	7.7	0.2	0.2	0.2
Phosphorus, total (as P)	0.014	0.046	0.013	0.011	0.048	0.013	0.009	0.012	0.020	0.039
Phosphorus, ortho, dissolved (as P)	<0.002	--	--	--	--	--	--	--	--	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	0.918	--	--	--	--	--	--	--	--	--
Nitrogen, ammonia, dissolved (as N)	<0.015	--	--	--	--	--	--	--	--	--
Nitrogen, amm. + diss., total (as N)	0.72	--	--	--	--	--	--	--	--	--

7-14-04




8-17-04

DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER






9-23-04




WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS

pH IN STANDARD UNITS



SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS



430551088273500 OCONOMOWOC LAKE NO. 1 (CENTER) AT OCONOMOWOC, WI


LOCATION.--Lat 43°05'51", long 88°27'35", in NW 1/4 SE 1/4 sec.2, T.7 N., R.17 E., Waukesha County, Hydrologic Unit 07090001, at Oconomowoc.


PERIOD OF RECORD.--March 1986 to current year.

REMARKS.--Lake sampled near center at the deep hole. Lake ice-covered during February sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, FEBRUARY 19 TO AUGUST 17, 2004
(Milligrams per liter unless otherwise indicated)

Date	Feb-19	Apr-13	Jun-8	Jul-15	Aug-17
Lake stage (ft)	7.09	7.82	8.23	7.11	7.03
Secchi depth (m)	--	2.7	6.8	5.0	3.3
Depth of sample (m)	0.5	18	0.5	18	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	--	6.3	2.1	--
Water temperature ($^{\circ}\text{C}$)	0.3	2.8	7.2	6.2	22.2
Specific conductance ($\mu\text{S/cm}$)	642	661	555	555	569
pH	7.9	7.4	8.3	8.1	7.7
Dissolved oxygen (mg/L)	14.2	6.7	12.9	11.8	10.0
Phosphorus, total (as P)	0.010	0.016	0.015	0.014	0.022
Phosphorus, ortho, dissolved (as P)	--	--	<0.002	--	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	--	--	0.158	--	--
Nitrogen, ammonia, dissolved (as N)	--	--	0.028	--	--
Nitrogen, amm. + org., total (as N)	--	--	0.56	--	--
Nitrogen, total (as N)	--	--	0.72	--	--
Color (Pt-Co. scale)	--	--	10	--	--
Turbidity (NTU)	--	--	1.1	--	--
Hardness, as CaCO_3	--	--	250	--	--
Calcium, dissolved (Ca)	--	--	44.1	--	--
Magnesium, dissolved (Mg)	--	--	34.8	--	--
Sodium, dissolved (Na)	--	--	19.5	--	--
Potassium, dissolved (K)	--	--	2	--	--
Alkalinity, as CaCO_3	--	--	206	--	--
Sulfate, dissolved (SO_4)	--	--	26.3	--	--
Chloride, dissolved (Cl)	--	--	43.9	--	--
Silica, dissolved (SiO_2)	--	--	5.26	--	--
Solids, dissolved, at 180°C	--	--	312	--	--
Iron, dissolved (Fe) ($\mu\text{g/L}$)	--	--	<100	--	--
Manganese, dissolved, (Mn) ($\mu\text{g/L}$)	--	--	<1	--	--





Surface total phosphorus, chlorophyll a concentrations, Secchi depths,
and TSI data for Oconomowoc Lake, Center Site, at Oconomowoc, Wisconsin.

(Circles on the first two plots indicate laboratory detection limit for selected analyses.
Actual concentrations for these particular analyses are less than the plotted circles.)

430609088262200 OCONOMOWOC LAKE NO. 2 (OFF HEWITT POINT) AT OCONOMOWOC, WI


LOCATION.--Lat 43°06'09", long 88°26'22", in NW 1/4 NW 1/4 sec.1, T.7 N., R.17 E., Waukesha County, Hydrologic Unit 07090001, at Oconomowoc.


PERIOD OF RECORD.--March 1986 to current year.

REMARKS.--Lake sampled at the deepest point in northeast bay near Hewitt Point. Lake ice-covered during February sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, FEBRUARY 19 TO AUGUST 17, 2004
(Milligrams per liter unless otherwise indicated)

Date	Feb-19	Apr-13	Jun-8	Jul-15	Aug-17
Lake stage (ft)	7.09	7.82	8.23	7.11	7.03
Secchi depth (m)	--	6.2	7.6	4.8	3.1
Depth of sample (m)	0.5	14	0.5	14.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	--	1.1	--	2.1
Water temperature ($^{\circ}\text{C}$)	1.0	4.2	7.9	11.4	21.9
Specific conductance ($\mu\text{S/cm}$)	660	711	602	557	628
pH	7.8	7.3	8.2	8.2	8.3
Dissolved oxygen (mg/L)	13.3	6.1	12.2	9.9	11.9
Phosphorus, total (as P)	0.008	0.009	0.012	0.013	0.010





Surface total phosphorus, chlorophyll a concentrations, Secchi depths,
and TSI data for Oconomowoc Lake, Hewitt Point, at Oconomowoc, Wisconsin.

(Circles on the first two plots indicate laboratory detection limit for selected analyses.
Actual concentrations for these particular analyses are less than the plotted circles.)

430723088252100 OKAUCHEE LAKE AT OKAUCHEE, WI

LOCATION.--Lat 43°07'23", long 88°25'21", in SE 1/4 SE 1/4, sec.25, T.8 N., R.17 E., Waukesha County, Hydrologic Unit 07090001, at Okauchee.

DRAINAGE AREA.--80.7 mi².

PERIOD OF RECORD.--February 1984 to current year.

LAKE-STAGE GAGE.--Datum of gage is 869.00 ft above sea level.

REMARKS.--Lake sampled near center at the deep hole. Lake ice-covered during February sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, FEBRUARY 19 TO AUGUST 17, 2004

(Milligrams per liter unless otherwise indicated)

Date	Feb-19	Apr-13	Jun-8	Jul-15	Aug-17
Lake stage (ft)	3.88	4.72	4.70	4.78	4.70
Secchi depth (m)	--	1.7	2.6	2.0	2.0
Depth of sample (m)	0.5	27.5	0.5	27.5	0.5
Chlorophyll a, phytoplankton (µg/L)	--	--	11.9	6.3	--
Water temperature (°C)	0.4	2.2	6.6	5.6	21.7
Specific conductance (µS/cm)	637	644	555	552	557
pH	8.1	7.7	8.3	8.1	7.7
Dissolved oxygen (mg/L)	13.7	4.1	13.5	12.3	10.4
Phosphorus, total (as P)	0.014	0.014	0.027	0.020	0.027
Phosphorus, ortho, dissolved (as P)	--	--	<0.002	--	--
Nitrogen, NO ₂ + NO ₃ , diss. (as N)	--	--	0.261	--	0.147
Nitrogen, ammonia, dissolved (as N)	--	--	0.002	--	0.028
Nitrogen, amm. + diss., total (as N)	--	--	--	--	0.75
Nitrogen, amm. + org., total (as N)	--	--	0.6	--	--
Nitrogen, total (as N)	--	--	0.86	--	--
Color (Pt-Co. scale)	--	--	10	--	--
Turbidity (NTU)	--	--	2.7	--	--
Hardness, as CaCO ₃	--	--	260	--	--
Calcium, dissolved (Ca)	--	--	47.9	--	--
Magnesium, dissolved (Mg)	--	--	34.2	--	--
Sodium, dissolved (Na)	--	--	15.5	--	--
Potassium, dissolved (K)	--	--	2	--	--
Alkalinity, as CaCO ₃	--	--	216	--	--
Sulfate, dissolved (SO ₄)	--	--	26.9	--	--
Chloride, dissolved (Cl)	--	--	37.5	--	--
Silica, dissolved (SiO ₂)	--	--	2.74	--	--
Solids, dissolved, at 180°C	--	--	318	--	--
Iron, dissolved (Fe) (µg/L)	--	--	<100	--	--
Manganese, dissolved, (Mn) (µg/L)	--	--	<1	--	--

2-19-04


4-13-04

6-8-04

7-15-04


8-17-04

DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER




WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS

PH IN STANDARD UNITS



SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS



Surface total phosphorus, chlorophyll a concentrations, Secchi depths,
and TSI data for Okauchee Lake, near Okauchee, Wisconsin.

(Circles on the first two plots indicate laboratory detection limit for selected analyses.
Actual concentrations for these particular analyses are less than the plotted circles.)

430759088244200 OKAUCHEE LAKE, NO. 1, NEAR OKAUCHEE, WI

LOCATION.--Lat 43°07'59", long 88°24'42", in NE 1/4 NW 1/4 sec.30, T.8 N., R.18 E., Waukesha County, Hydrologic Unit 07090001, near Okauchee.


PERIOD OF RECORD.--April 1986 to current year.

LAKE-STAGE GAGE.--Datum of gage is 869.00 ft above sea level.

REMARKS.--Lake sampled in Crane's Nest Bay, in the northeast part of the lake, at an approximate depth of 2 m. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, APRIL 13 TO AUGUST 17, 2004
(Milligrams per liter unless otherwise indicated)

Date	<u>Apr-13</u>	<u>Jun-8</u>	<u>Jul-15</u>	<u>Aug-17</u>
Lake stage (ft)	4.72	4.70	4.78	4.70
Secchi depth (m)	1.8	1.4	1.1	1.3
Depth of sample (m)	0.5	0.5	0.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	9.2	21.9	10.1	14.2
Water temperature ($^{\circ}\text{C}$)	6.1	24.7	25.2	21.7
Specific conductance ($\mu\text{S/cm}$)	552	575	592	565
pH	8.4	8.4	7.9	8.2
Dissolved oxygen (mg/L)	13.3	11.1	7.6	11.2
Phosphorus, total (as P)	0.022	0.034	0.041	0.026



Surface total phosphorus, chlorophyll a concentrations, Secchi depths,
and TSI data for Okauchee Lake, No. 1, near Okauchee, Wisconsin.

(Triangles in Secchi plot indicate maximum depth at sampling site.
Actual Secchi depth on these days was greater than the plotted triangles.)

430645088264500 OKAUCHEE LAKE, NO. 2, AT OKAUCHEE, WI

LOCATION.--Lat 43°06'45", long 88°26'45", in SE 1/4 NE 1/4 sec.35, T.8 N., R.17 E., Waukesha County, Hydrologic Unit 07090001, at Okauchee.


PERIOD OF RECORD.--April 1986 to current year.

LAKE-STAGE GAGE.--Datum of gage is 869.00 ft above sea level.

REMARKS.--Lake sampled in Lower Okauchee Lake, at an approximate depth of 5 m. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, APRIL 13 TO AUGUST 17, 2004
(Milligrams per liter unless otherwise indicated)

Date	<u>Apr-13</u>	<u>Jun-8</u>	<u>Jul-15</u>	<u>Aug-17</u>
Lake stage (ft)	4.72	4.70	4.78	4.70
Secchi depth (m)	1.0	1.9	2.3	1.7
Depth of sample (m)	0.5	0.5	0.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	10.1	6.5	3.7	14.7
Water temperature ($^{\circ}\text{C}$)	8.2	23.9	25.5	21.8
Specific conductance ($\mu\text{S/cm}$)	544	549	534	533
pH	8.5	8.3	8.1	8.3
Dissolved oxygen (mg/L)	13.1	11.2	10.0	13.4
Phosphorus, total (as P)	0.027	0.019	0.020	0.026



Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Okauchee Lake, No. 2, near Okauchee, Wisconsin.

(Circles on the first two plots indicate laboratory detection limit for selected analyses.
Actual concentrations for these particular analyses are less than the plotted circles.)

(Triangles in Secchi plot indicate maximum depth at sampling site.
Actual Secchi depth on these days was greater than the plotted triangles.)

430642088252400 OKAUCHEE LAKE, NO. 3, AT OKAUCHEE, WI

LOCATION.--Lat 43°06'42", long 88°25'24", in NE 1/4 SE 1/4 sec.36, T.8 N., R.17 E., Waukesha County, Hydrologic Unit 07090001, at Okauchee.


PERIOD OF RECORD.--April 1986 to current year.

LAKE-STAGE GAGE.--Datum of gage is 869.00 ft above sea level.

REMARKS.--Lake sampled in Ice House Bay, in the southern part of the lake, at an approximate depth of 4 m. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, APRIL 13 TO AUGUST 17, 2004
(Milligrams per liter unless otherwise indicated)

Date	<u>Apr-13</u>	<u>Jun-8</u>	<u>Jul-15</u>	<u>Aug-17</u>
Lake stage (ft)	4.72	4.70	4.78	4.7
Secchi depth (m)	1.2	1.7	1.4	--
Depth of sample (m)	0.5	0.5	0.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	11.1	6.9	9.7	12.4
Water temperature ($^{\circ}\text{C}$)	7.0	22.0	25.1	21.3
Specific conductance ($\mu\text{S/cm}$)	553	545	540	538
pH	8.5	8.3	8.1	8.3
Dissolved oxygen (mg/L)	13.2	11.1	9.5	11.7
Phosphorus, total (as P)	0.023	0.023	0.022	0.021



Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Okauchee Lake, No. 3, near Okauchee, Wisconsin.

(Triangles in Secchi plot indicate maximum depth at sampling site.
Actual Secchi depth on these days was greater than the plotted triangles.)

430757088261700 OKAUCHEE LAKE, NO. 4, AT OKAUCHEE, WI

LOCATION.--Lat 43°07'57", long 88°26'17", in NW 1/4 NW 1/4 sec.25, T.8 N., R.17 E., Waukesha County, Hydrologic Unit 07090001, at Okauchee.


PERIOD OF RECORD.--June 1986 to current year.

LAKE-STAGE GAGE.--Datum of gage is 869.00 ft above sea level.

REMARKS.--Lake sampled near McDowell (Crazyman's) Island, in the northwest bay of the lake, at an approximate depth of 2 m. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, APRIL 13 TO AUGUST 17, 2004
(Milligrams per liter unless otherwise indicated)

Date	<u>Apr-13</u>	<u>Jun-8</u>	<u>Jul-15</u>	<u>Aug-17</u>
Lake stage (ft)	4.72	4.70	4.78	4.70
Secchi depth (m)	1.4	1.4	2.0	1.6
Depth of sample (m)	0.5	0.5	0.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	8.7	4.5	5.1	5.7
Water temperature ($^{\circ}\text{C}$)	6.4	23.2	24.2	21.5
Specific conductance ($\mu\text{S/cm}$)	546	552	538	531
pH	8.4	8.4	8.2	8.3
Dissolved oxygen (mg/L)	13.7	10.3	10.1	13.4
Phosphorus, total (as P)	0.023	0.015	0.022	0.016



Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Okauchee Lake, No. 4, near Okauchee, Wisconsin.

(Circles on the first two plots indicate laboratory detection limit for selected analyses.
Actual concentrations for these particular analyses are less than the plotted circles.)

(Triangles in Secchi plot indicate maximum depth at sampling site.
Actual Secchi depth on these days was greater than the plotted triangles.)

424905088204000 POTTER LAKE NEAR MUKWONAGO, WI

LOCATION.--Lat 42°49'05", long 88°20'40", in NW 1/4 SW 1/4 sec.11, T.4 N., R.18 E., Walworth County, Hydrologic Unit 07120006, 3.3 mi south of Mukwonago.

PERIOD OF RECORD.--February 1993 to current year.

REMARKS.--Lake sampled at the deep hole. Lake ice-covered during February sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, APRIL 8 TO JULY 27, 2004
(Milligrams per liter unless otherwise indicated)

Date	Apr-8	Jun-7	Jul-12	Jul-27
Lake stage (ft)	6.74	8.44	8.36	8.08
Secchi depth (m)	2.3	2.5	1.7	1.3
Depth of sample (m)	0.5	7	0.5	7
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	7.6	--	3.1	--
Water temperature ($^{\circ}\text{C}$)	10.6	10.5	23.3	15.7
Specific conductance ($\mu\text{S/cm}$)	404	404	387	550
pH	8.8	8.8	8.5	7.3
Dissolved oxygen (mg/L)	12.0	11.7	10.4	0.4
Phosphorus, total (as P)	0.024	0.020	0.038	0.134
Phosphorus, ortho, dissolved (as P)	0.005	--	--	0.018
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	0.019	--	--	<0.019
Nitrogen, ammonia, dissolved (as N)	0.026	--	--	0.024
Nitrogen, amm. + diss., total (as N)	--	--	--	0.89
Nitrogen, amm. + org., total (as N)	1.2	--	--	--
Color (Pt-Co. scale)	10	--	--	--
Turbidity (NTU)	1.9	--	--	--
Hardness, as CaCO_3	150	--	--	--
Calcium, dissolved (Ca)	25.5	--	--	--
Magnesium, dissolved (Mg)	20.4	--	--	--
Sodium, dissolved (Na)	23.1	--	--	--
Potassium, dissolved (K)	2	--	--	--
Alkalinity, as CaCO_3	121	--	--	--
Sulfate, dissolved (SO_4)	7	--	--	--
Chloride, dissolved (Cl)	49.8	--	--	--
Silica, dissolved (SiO_2)	0.066	--	--	--
Solids, dissolved, at 180°C	234	--	--	--
Iron, dissolved (Fe) ($\mu\text{g/L}$)	<100	--	--	--
Manganese, dissolved, (Mn) ($\mu\text{g/L}$)	<1	--	--	--





4-8-04

6-7-04





7-12-04

7-27-04

DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER



WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS





SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS

424905088204000 POTTER LAKE NEAR MUKWONAGO, WI--CONTINUED

WATER-QUALITY DATA, AUGUST 9 TO SEPTEMBER 23, 2004
 (Milligrams per liter unless otherwise indicated)

Date	<u>Aug-9</u>			<u>Aug-24</u>			<u>Sep-23</u>
Lake stage (ft)	8.06			7.89			7.62
Secchi depth (m)	1.3			1.3			0.7
Depth of sample (m)	0.5	5	7	0.5	7	0.5	7
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	18.3	--	--	27.1	--	13.5	--
Water temperature ($^{\circ}\text{C}$)	24.3	19.5	16.0	22.4	17.7	21.9	17.8
Specific conductance ($\mu\text{S/cm}$)	455	507	575	471	543	485	595
pH	7.8	7.8	7.7	8.0	6.8	8.1	6.8
Dissolved oxygen (mg/L)	8.3	0.2	0.1	9.3	0.1	9.1	0.1
Phosphorus, total (as P)	0.049	0.064	0.162	0.054	0.124	0.047	0.154





423246088175800 POWERS LAKE AT POWERS LAKE, WI

LOCATION.--Lat 42°32'46", long 88°17'58", in NW 1/4 SE 1/4 sec.13, T.1 N., R.18 E., Walworth County, Hydrologic Unit 07120006, at Powers Lake.

DRAINAGE AREA.--3.42 mi².


PERIOD OF RECORD.--March 1986 to August 1996, and April 1998 to current year.

REMARKS.--Lake sampled near center at the deep hole. Lake ice-covered during February sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, FEBRUARY 25 TO JUNE 7, 2004
(Milligrams per liter unless otherwise indicated)


Date	Feb-25	Apr-8	Jun-7
Lake stage (ft)	9.83	10.36	10.39
Secchi depth (m)	--	4.4	4.6
Depth of sample (m)	0.5	9.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	--	3.5
Water temperature (°C)	1.0	3.6	21.4
Specific conductance ($\mu\text{S/cm}$)	543	594	491
pH	8.1	7.6	8.2
Dissolved oxygen (mg/L)	14.8	5.8	11.2
Phosphorus, total (as P)	--	--	0.017
Phosphorus, ortho, dissolved (as P)	--	<0.002	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	--	<0.019	--
Nitrogen, ammonia, dissolved (as N)	--	0.009	--
Nitrogen, amm. + org., total (as N)	--	0.6	--
Color (Pt-Co. scale)	--	--	--
Turbidity (NTU)	--	--	--
Hardness, as CaCO_3	--	--	--
Calcium, dissolved (Ca)	--	--	--
Magnesium, dissolved (Mg)	--	--	--
Sodium, dissolved (Na)	--	--	--
Potassium, dissolved (K)	--	--	--
Alkalinity, as CaCO_3	--	--	--
Sulfate, dissolved (SO_4)	--	--	--
Chloride, dissolved (Cl)	--	--	--
Silica, dissolved (SiO_2)	--	--	--
Solids, dissolved, at 180°C	--	--	--
Iron, dissolved (Fe) ($\mu\text{g/L}$)	--	<100	--
Manganese, dissolved, (Mn) ($\mu\text{g/L}$)	--	<1	--

2-25-04




4-8-04


DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER




6-7-04



WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS



PH IN STANDARD UNITS





SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS

423246088175800 POWERS LAKE AT POWERS LAKE, WI--CONTINUED

WATER-QUALITY DATA, JULY 12 TO SEPTEMBER 23, 2004
(Milligrams per liter unless otherwise indicated)

	<u>Jul-12</u>	<u>Aug-16</u>	<u>Sep-23</u>
Date	10.09	9.88	9.92
Lake stage (ft)	2.8	3.1	1.4
Secchi-depth (m)			
Depth of sample (m)	0.5	9.5	8
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	3.2	--	4.1
Water temperature ($^{\circ}\text{C}$)	24.1	16.8	20.4
Specific conductance ($\mu\text{S/cm}$)	493	503	507
pH	8.1	7.3	8.1
Dissolved oxygen (mg/L)	11.6	0.3	5.9
Phosphorus, total (as P)	0.016	0.024	0.015
		0.054	0.020
			0.020
			0.047





Surface total phosphorus, chlorophyll a concentrations, Secchi depths,
and TSI data for Powers Lake, at Powers Lake, Wisconsin.

(Circles on the first two plots indicate laboratory detection limit for selected analyses.
Actual concentrations for these particular analyses are less than the plotted circles.)

455034091493300 SPOONER LAKE, AT DEEP HOLE, NEAR SPOONER, WI

LOCATION.--Lat 45°50'34", long 91°49'33", in NE 1/4 NE 1/4 sec.27, T.39 N., R.12 W., Washburn County, Hydrologic Unit 07030001, near Spooner.

PERIOD OF RECORD.--June 2002 to current year.

REMARKS.--Lake sampled at deepest hole. Lake ice-covered during March sampling. Water-quality analyses by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, JUNE 9 TO JULY 12, 2004
(Milligrams per liter unless otherwise indicated)

Date	<u>Jun-9</u>	<u>Jun-21</u>		<u>Jul-2</u>		<u>Jul-12</u>	
Lake stage (ft)	7.08			7.08		--	
Secchi depth (m)	1.9		2.5	1.7		2.2	
Depth of sample (m)	0.5	4.5	0.5	4	0.5	4.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	9.1	--	--	6.2	--	8.2	--
Water temperature ($^{\circ}\text{C}$)	22.0	21.0	20.4	20.1	21.5	19.2	23.1
Specific conductance ($\mu\text{S/cm}$)	179	185	184	184	184	193	176
pH	8.0	7.6	8.6	8.5	8.4	7.8	8.8
Dissolved oxygen (mg/L)	9.1	5.5	8.9	8.6	10.8	4.2	10.0
Phosphorus, total (as P)	0.035	0.031	0.035	0.043	0.024	0.033	0.023
							0.027





6-9-04

6-21-04

7-2-04





7-12-04

DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER



WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS

PH IN STANDARD UNITS



SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS

455034091493300 SPOONER LAKE, AT DEEP HOLE, NEAR SPOONER, WI--CONTINUED

WATER-QUALITY DATA, JULY 20 TO AUGUST 25, 2004
 (Milligrams per liter unless otherwise indicated)

Date	<u>Jul-20</u>	<u>Jul-28</u>	<u>Aug-12</u>	<u>Aug-25</u>
Lake stage (ft)	7.02	6.95	7.12	6.40
Secchi depth (m)	1.9	1.7	3.0	2.5
Depth of sample (m)	0.5	4.5	0.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	8.1	--	8.5	6.7
Water temperature ($^{\circ}\text{C}$)	25.8	23.5	24.5	19.9
Specific conductance ($\mu\text{S/cm}$)	169	187	169	151
pH	8.8	8.0	8.7	9.0
Dissolved oxygen (mg/L)	9.9	0.7	8.8	8.8
Phosphorus, total (as P)	0.025	0.031	0.033	0.026
Phosphorus, ortho, dissolved (as P)	--	--	0.003	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	--	--	<0.019	--
Nitrogen, ammonia, dissolved (as N)	--	--	<0.015	--
Nitrogen, amm. + diss., total (as N)	--	--	0.52	--





7-20-04

7-28-04





8-12-04

8-25-04





DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER




WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS



PH IN STANDARD UNITS



SPECIFIC CONDUCTANCE (S.C.) IN MICROSIEMENS PER CENTIMETER AT 25 DEGREES CELSIUS



Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Spooner Lake, Deep Hole, near Spooner, Wisconsin.

454945091483900 SPOONER LAKE, SOUTHEAST SITE, NEAR SPOONER, WI

LOCATION.--Lat 45°49'45", long 91°48'39", in SW 1/4 SE 1/4 sec.26, T.39 N., R.12 W., Washburn County, Hydrologic Unit 07030001, near Spooner.

PERIOD OF RECORD.--June 2002 to current year.





REMARKS.--Lake sampled at deepest hole. Lake ice-covered during March sampling. Water-quality analyses by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, JUNE 9 TO JULY 12, 2004





(Milligrams per liter unless otherwise indicated)

Date	<u>Jun-9</u>	<u>Jun-21</u>	<u>Jul-2</u>	<u>Jul-12</u>
Lake stage (ft)	7.08	6.90	7.08	7.10
Secchi depth (m)	1.9	2.3	2.4	2.0
Depth of sample (m)	0.5	2	0.5	2
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	7.6	--	8.1	12.2
Water temperature ($^{\circ}\text{C}$)	22.2	22.2	20.5	23.5
Specific conductance ($\mu\text{S/cm}$)	177	177	190	194
pH	8.5	8.5	8.3	8.4
Dissolved oxygen (mg/L)	9.9	10.1	8.4	9.6
Phosphorus, total (as P)	0.037	0.036	0.058	0.040
			0.050	0.034
			0.033	0.040
			0.037	





DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER



WATER TEMPERATURE (W.T.) IN DEGREES CELSIUS



PH IN STANDARD UNITS





454945091483900 SPOONER LAKE, SOUTHEAST SITE, NEAR SPOONER, WI CONTINUED

454945091483900 SPOONER LAKE, SOUTHEAST SITE, NEAR SPOONER, WI--CONTINUED

WATER-QUALITY DATA, JULY 20 TO AUGUST 25, 2004
 (Milligrams per liter unless otherwise indicated)

Date	Jul-20	Jul-28	Aug-12	Aug-25
Lake stage (ft)	7.02	6.95	7.12	6.94
Secchi depth (m)	1.6	1.6	2.8	1.9
Depth of sample (m)	0.5	2	0.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	16.5	--	19.6	--
Water temperature ($^{\circ}\text{C}$)	26.0	23.7	23.6	20.2
Specific conductance ($\mu\text{S/cm}$)	197	206	201	200
pH	8.5	8.0	8.4	8.3
Dissolved oxygen (mg/L)	10.0	8.3	9.1	9.6
Phosphorus, total (as P)	0.043	0.044	0.041	0.034
			0.04	0.032





Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Spooner Lake, Southeast Site, near Spooner, Wisconsin.

424854088123300 TICHIGAN LAKE NEAR WATERFORD, WI


LOCATION.--Lat 42°48'54", long 88°12'33", in se 1/4 SE 1/4 sec.11, T.4 N., R.19 E., Racine County, Hydrologic Unit 07120006, 3.5 mi north of Waterford.

PERIOD OF RECORD.--March 1994 to August 1996, April 2003 to current year.

REMARKS.--Lake sampled near center at the deep hole. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.


WATER-QUALITY DATA, FEBRUARY 18 TO JUNE 7, 2004
(Milligrams per liter unless otherwise indicated)


Date	<u>Feb-18</u>	<u>Apr-12</u>	<u>Jun-7</u>
Lake stage (ft)	1.49	1.48	1.34
Secchi depth (m)	--	1.3	1.3
Depth of sample (m)	0.5	0.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	--	19.9
Water temperature ($^{\circ}\text{C}$)	0.4	2.4	5.9
Specific conductance ($\mu\text{S/cm}$)	1140	1220	1010
pH	7.5	7.1	7.7
Dissolved oxygen (mg/L)	13.3	7.0	5.6
Phosphorus, total (as P)	0.082	0.121	0.137
			0.04
			0.376



WATER-QUALITY DATA, JUNE 12 TO AUGUST 23, 2004
 (Milligrams per liter unless otherwise indicated)

Date	Jun-12	Jul-16	Aug-23
Lake stage (ft)	1.62	1.51	1.30
Secchi depth (m)	1.9	1.1	0.7
Depth of sample (m)	0.5	18.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	11.6	--	14.6
Water temperature ($^{\circ}\text{C}$)	24.9	9.9	22.3
Specific conductance ($\mu\text{S/cm}$)	800	1050	874
pH	8.4	7.3	8.5
Dissolved oxygen (mg/L)	11.9	0.2	0.2
Phosphorus, total (as P)	0.037	0.609	0.054
		0.038	0.804
			1.080





435430089350700 TWIN LAKES, EAST TWIN LAKE, NEAR WESTFIELD, WI


LOCATION.--Lat 43°54'30", long 89°35'07", in NE 1/4 NE 1/4 sec.3, T.17 N., R.8 E., Marquette County, Hydrologic Unit 04030201, near Westfield.


PERIOD OF RECORD.--June 2002 current year.

REMARKS.--Lake sampled at deepest hole. Water-quality analyses by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, APRIL 20 TO AUGUST 23, 2004
(Milligrams per liter unless otherwise indicated)

Date	Apr-20	Jun-22	Jul-8	Aug-23
Lake stage (ft)	9.02	9.73	9.83	10.05
Secchi depth (m)	1.5	1.0	0.8	0.8
Depth of sample (m)	0.5	0.5	0.5	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	12.2	12.0	14.7	14.0
Water temperature ($^{\circ}\text{C}$)	14.1	22.1	20.0	20.9
Specific conductance ($\mu\text{S/cm}$)	221	168	192	218
pH	8.3	8.7	8.4	8.5
Dissolved oxygen (mg/L)	10.9	9.2	7.7	10.4
Phosphorus, total (as P)	0.034	0.040	0.052	0.051





Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Twin Lakes, East Twin Lake, near Westfield, Wisconsin.

435438089352300 TWIN LAKES, WEST TWIN LAKE, NEAR WESTFIELD, WI


LOCATION.--Lat 43°54'38", long 89°35'23", in SW 1/4 SE 1/4 sec.30, T.17 N., R.8 E., Marquette County, Hydrologic Unit 04030201, near Westfield.


PERIOD OF RECORD.--June 2002 to current year.

REMARKS.--Lake sampled at deepest hole. Water-quality analyses by Wisconsin State Laboratory of Hygiene.

**WATER-QUALITY DATA, APRIL 20 TO AUGUST 23, 2004
(Milligrams per liter unless otherwise indicated)**

Date	Apr-20	Jun-22	Jul-8	Aug-23
Lake stage (ft)	9.02	9.73	9.83	10.05
Secchi depth (m)	1.8	1.0	1.1	1.2
Depth of sample (m)	0.5	8	0.5	8.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	8.4	--	4.9	--
Water temperature ($^{\circ}\text{C}$)	13.4	6.0	21.3	6.8
Specific conductance ($\mu\text{S/cm}$)	299	449	201	564
pH	8.2	6.8	8.8	7.1
Dissolved oxygen (mg/L)	11.9	0.3	10.9	0.3
Phosphorus, total (as P)	0.040	0.230	0.022	0.259
			0.030	0.607
			0.018	0.019
			0.102	





Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Twin Lakes, West Twin Lake, near Westfield, Wisconsin.

424614088123600 WATERFORD WATERWAY, FOX RIVER SITE, NEAR WATERFORD, WI

LOCATION.--Lat 42°46'14", long 88°12'36", in NE 1/4 NE 1/4 NE 1/4 sec.35, T.4 N., R.19 E., Racine County, Hydrologic Unit 07120006, near Waterford.


PERIOD OF RECORD.--April 2003 to current year.

REMARKS.--Lake sampled at Fox River site. River was ice-covered during February sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, FEBRUARY 18 TO JUNE 7, 2004

(Milligrams per liter unless otherwise indicated)


Date	Feb-18	Apr-12	Jun-7
Lake stage (ft)	1.49	1.48	1.34
Secchi depth (m)	--	0.5	0.7
Depth of sample (m)	0.5	0.5	0.25
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	60.1	39.9
Water temperature (°C)	0.1	8.4	25.1
Specific conductance ($\mu\text{S/cm}$)	1370	1020	813
pH	7.6	8.6	7.8
Dissolved oxygen (mg/L)	16.4	16.1	9.6
Phosphorus, total (as P)	0.034	0.076	0.080




424614088123600 WATERFORD WATERWAY, FOX RIVER SITE, NEAR WATERFORD, WI--CONTINUED

WATER-QUALITY DATA, JULY 12 TO SEPTEMBER 23, 2004
 (Milligrams per liter unless otherwise indicated)

Date	<u>Jul-12</u>	<u>Aug-16</u>	<u>Sep-23</u>
Lake stage (ft)	1.62	1.51	1.30
Secchi depth (m)	0.4	0.2	0.1
Depth of sample (m)	0.25	0.5	0.25
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	144.0	149.0	97.1
Water temperature ($^{\circ}\text{C}$)	26.3	22.0	22.2
Specific conductance ($\mu\text{S/cm}$)	869	904	995
pH	8.3	8.5	8.8
Dissolved oxygen (mg/L)	20.0	16.6	14.3
Phosphorus, total (as P)	0.140	0.166	0.179





Surface total phosphorus, chlorophyll a concentrations, Secchi depths, and TSI data for Waterford Waterway, Fox River Site, near Waterford, Wisconsin.

441730090432900 WAZEE LAKE AT DEEP HOLE NEAR BLACK RIVER FALLS, WI

LOCATION.--Lat 44°17'30", long 90°43'29", in SE 1/4 SE 1/4 sec.15, T.21 N., R.3 W., Jackson County, Hydrologic Unit 07040007, 7.0 mi east of Black River Falls.


PERIOD OF RECORD.--November 1999 to current year.

REMARKS.--Date published under station number 441721090431700 in water year 2000. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.


WATER-QUALITY DATA, DECEMBER 6, 2003
(Milligrams per liter unless otherwise indicated)


Date	Dec-6
Lake stage (ft)	3.42
Secchi depth (m)	12.8
Depth of sample (m)	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	6.79
Water temperature ($^{\circ}\text{C}$)	5.4
Specific conductance ($\mu\text{S/cm}$)	246
pH	8.0
Dissolved oxygen (mg/L)	12.7
Phosphorus, total (as P)	0.006

12-6-03
DISSOLVED OXYGEN (D.O.) IN MILLIGRAMS PER LITER



PH IN STANDARD UNITS





Surface total phosphorus, chlorophyll a concentrations, Secchi depths,
and TSI data for Wazee Lake, Deep Hole, near Black River Falls, Wisconsin.

(Circles on the first two plots indicate laboratory detection limit for selected analyses.
Actual concentrations for these particular analyses are less than the plotted circles.)

424848088083100 WIND LAKE AT OUTLET AT WIND LAKE, WI

LOCATION.--Lat 42°48'48" long 88°08'31", in NE 1/4 NW 1/4 sec.16, T.4 N., R.20 E., Racine County, Hydrologic Unit 07120006, at Wind Lake.

DRAINAGE AREA.--39.6 mi².

PERIOD OF RECORD.--March 1985 to current year. Prior to October 2000, published as "Wind Lake Outlet".

REVISED RECORDS.--WDR WI-91-1: 1988(m).

GAGE.--Water-stage recorder and concrete dam. Datum of gage is 760.30 ft above sea level. Prior to Oct. 2, 1987, nonrecording gage at same site and datum.


REMARKS.--Lake level regulated by dam with two 10-foot gates at outlet. Lake ice-covered Dec. 1 to Mar. 11. Prior to October 1987, published as Wind Lake at Wind Lake, Wis. Gage-height telemeter at station.

EXTREMES FOR PERIOD OF RECORD.--Maximum gage height, 8.93 ft, June 15, 1999; minimum recorded, 5.95 ft, Jan. 2, 1996.

EXTREMES FOR CURRENT YEAR.--Maximum recorded gage height, 8.57 ft, June 7; minimum recorded, 7.43 ft, Oct. 28.

GAGE HEIGHT, FEET
WATER YEAR OCTOBER 2003 TO SEPTEMBER 2004
DAILY MEAN VALUES

DAY	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP
1	7.55	7.47	7.84	8.15	8.14	8.18	8.14	8.23	8.18	8.11	8.18	8.07
2	7.53	7.54	7.84	8.16	8.13	8.18	8.11	8.20	8.12	8.11	8.17	8.06
3	7.52	7.63	7.84	8.17	8.14	8.15	8.13	8.16	8.10	8.13	8.17	8.06
4	7.53	7.69	7.85	8.16	8.13	8.12	8.13	8.12	8.14	8.20	8.23	8.05
5	7.51	7.71	7.88	8.16	8.12	8.26	8.11	8.13	8.33	8.23	8.20	8.04
6	7.51	7.71	7.89	8.13	8.13	8.32	8.11	8.14	8.49	8.25	8.17	8.03
7	7.50	7.70	7.90	8.11	8.12	8.31	8.13	8.15	8.52	8.30	8.15	8.02
8	7.50	7.68	7.92	8.08	8.11	8.24	8.21	8.14	8.35	8.32	8.14	8.01
9	7.50	7.67	7.95	8.07	8.10	8.16	8.22	8.22	8.15	8.35	8.13	7.99
10	7.50	7.67	8.13	8.06	8.08	8.08	8.20	8.20	8.15	8.27	8.11	7.97
11	7.49	7.67	8.17	8.06	8.08	8.03	8.17	8.19	8.22	8.23	8.09	7.95
12	7.50	7.67	8.20	8.07	8.08	8.01	8.13	8.16	8.26	8.25	8.07	7.94
13	7.49	7.66	8.25	8.09	8.07	8.07	8.08	8.24	8.29	8.25	8.06	7.93
14	7.53	7.64	8.30	8.10	8.07	8.15	8.05	8.24	8.31	8.22	8.04	7.91
15	7.52	7.64	8.33	8.12	8.06	8.23	8.08	8.12	8.28	8.21	8.03	7.90
16	7.51	7.64	8.28	8.13	8.05	8.22	8.15	8.04	8.22	8.19	8.01	7.91
17	7.49	7.64	8.21	8.14	8.04	8.11	8.22	8.10	8.29	8.20	8.05	7.89
18	7.48	7.73	8.20	8.15	8.04	8.07	8.22	8.32	8.27	8.18	8.05	7.87
19	7.48	7.75	8.17	8.15	8.04	8.10	8.23	8.31	8.18	8.18	8.04	7.85
20	7.47	7.74	8.14	8.15	8.05	8.17	8.19	8.15	8.16	8.18	8.03	7.84
21	7.48	7.76	8.13	8.15	8.07	8.21	8.21	8.10	8.26	8.20	8.01	7.82
22	7.47	7.77	8.16	8.15	8.06	8.19	8.18	8.27	8.33	8.25	7.99	7.81
23	7.46	7.79	8.20	8.16	8.08	8.12	8.16	8.42	8.29	8.25	7.99	7.80
24	7.46	7.78	8.21	8.18	8.08	8.05	8.17	8.52	8.28	8.23	7.99	7.78
25	7.50	7.78	8.19	8.19	8.09	8.00	8.20	8.53	8.25	8.22	8.04	7.77
26	7.49	7.79	8.17	8.20	8.10	8.06	8.22	8.48	8.33	8.24	8.04	7.76
27	7.48	7.80	8.14	8.23	8.11	8.09	8.23	8.39	8.40	8.25	8.05	7.75
28	7.47	7.83	8.14	8.22	8.13	8.14	8.23	8.26	8.43	8.25	8.10	7.74
29	7.47	7.82	8.15	8.20	8.14	8.23	8.22	8.18	8.34	8.23	8.11	7.71
30	7.47	7.83	8.15	8.18	---	8.21	8.23	8.19	8.19	8.21	8.09	7.69
31	7.47	---	8.15	8.16	---	8.20	---	8.23	---	8.20	8.08	---
MEAN	7.49	7.71	8.10	8.14	8.09	8.15	8.17	8.23	8.27	8.22	8.08	7.90
MAX	7.55	7.83	8.33	8.23	8.14	8.32	8.23	8.53	8.52	8.35	8.23	8.07
MIN	7.46	7.47	7.84	8.06	8.04	8.00	8.05	8.04	8.10	8.11	7.99	7.69



424915088083900 WIND LAKE AT WIND LAKE, WI


LOCATION.--Lat 42°49'15", long 88°08'39", in NW 1/4 SW 1/4 sec.9, T.4 N., R.20 E., Racine County, Hydrologic Unit 07120006, at Wind Lake.

PERIOD OF RECORD.--February 1985 to current year.

REMARKS.--Lake sampled near center at the deep hole. Lake ice-covered during February sampling. Water-quality analyses done by Wisconsin State Laboratory of Hygiene.

WATER-QUALITY DATA, FEBRUARY 18 TO JUNE 7, 2004
(Milligrams per liter unless otherwise indicated)


Date	Feb-18	Apr-12	Jun-7
Secchi depth (m)	--	2.5	2.1
Depth of sample (m)	0.5	15	0.5
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	--	2.5	7.0
Water temperature ($^{\circ}\text{C}$)	1.0	2.7	22.3
Specific conductance ($\mu\text{S/cm}$)	814	--	694
pH	7.9	7.2	7.9
Dissolved oxygen (mg/L)	13.1	4.1	9.2
Phosphorus, total (as P)	0.023	0.039	0.036
Phosphorus, ortho, dissolved (as P)	--	<0.002	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	--	0.513	--
Nitrogen, ammonia, dissolved (as N)	--	0.104	--
Nitrogen, ammn. + org., total (as N)	--	1.4	--
Nitrogen, total (as N)	--	1.9	--
Color (Pt-Co. scale)	--	30	--
Turbidity (NTU)	--	1.9	--
Hardness, as CaCO_3	--	250	--
Calcium, dissolved (Ca)	--	47	--
Magnesium, dissolved (Mg)	--	32.1	--
Sodium, dissolved (Na)	--	54.3	--
Potassium, dissolved (K)	--	3	--
Alkalinity, as CaCO_3	--	171	--
Sulfate, dissolved (SO_4)	--	46.9	--
Chloride, dissolved (Cl)	--	110	--
Silica, dissolved (SiO_2)	--	0.022	--
Solids, dissolved, at 180°C	--	446	--
Iron, dissolved (Fe) ($\mu\text{g/L}$)	--	<100	--
Manganese, dissolved, (Mn) ($\mu\text{g/L}$)	--	M	--




424915088083900 WIND LAKE AT WIND LAKE, WI--CONTINUED

WATER-QUALITY DATA, JULY 12 TO AUGUST 16, 2004
 (Milligrams per liter unless otherwise indicated)

Date	<u>Jul-12</u>			<u>Aug-16</u>			
Secchi depth (m)		1.1			2.3		
Depth of sample (m)	0.5	15.5	0.5	4	10	13	15
Chlorophyll a, phytoplankton ($\mu\text{g/L}$)	18.2	--	10.7	--	--	--	--
Water temperature ($^{\circ}\text{C}$)	25.6	14.5	21.3	21.0	18.2	14.9	14.7
Specific conductance ($\mu\text{S/cm}$)	672	764	688	689	712	766	774
pH	8.4	7.2	7.9	7.9	7.3	7.0	6.9
Dissolved oxygen (mg/L)	13.3	0.2	8.0	7.5	0.2	0.2	0.2
Phosphorus, total (as P)	0.035	0.360	0.033	0.027	0.082	0.375	0.436
Phosphorus, ortho, dissolved (as P)	<0.002	--	--	--	--	--	--
Nitrogen, $\text{NO}_2 + \text{NO}_3$, diss. (as N)	<0.019	--	--	--	--	--	--
Nitrogen, ammonia, dissolved (as N)	0.017	--	--	--	--	--	--
Nitrogen, amm. + diss., total (as N)	1.3	--	--	--	--	--	--





04082500 LAKE WINNEBAGO AT OSHKOSH, WI

LOCATION.--Lat 44°00'35", long 88°31'38", in NE 1/4 NE 1/4 sec.25, T.18 N., R.16 E., Winnebago County, Hydrologic Unit 04030203, at 905 Bay Shore Drive, 800 ft east of mouth of the upper Fox River.

DRAINAGE AREA.--5,880 mi², at lake outlet at Menasha Dam. Area of Lake Winnebago, 215 mi².

PERIOD OF RECORD.--October 1938 to current year in reports of Geological Survey. Records from July 1882 to September 1938 in files of Geological Survey and U.S. Army Corps of Engineers. A report on Fox River by U.S. Army Corps of Engineers, published as House Document No. 146, 67th Congress, 2nd session, contains semi-monthly records of inflow of Lake Winnebago for the period 1896-1917.

REVISED RECORD.--WDR WI-83-1: Drainage area.

GAGE.--Water-stage recorder. Nonrecording gage read once daily October 1938 to October 1978. Datum of gage is 745.05 ft above mean tide at New York City (levels by U.S. Army Corps of Engineers). Datum of Deuchman gage is 745.00 ft above mean tide at New York City.

REMARKS.--Lake elevations controlled by dams at Menasha and Neenah, which are operated in the interest of navigation. Crests of both dams are at elevation 746.73 ft. Present limits of regulation are from 21 1/4 in. above the crest of Menasha dam to crest during navigation season, plus additional 18 in. below crest during winter. Oshkosh staff gage gives true level of lake, while Deuchman gage readings are affected by loss of head in the channel between lake and dam. Data-collection platform and gage-height telemeter at station.

EXTREMES FOR PERIOD OF RECORD.--Maximum gage height observed, 5.33 ft (Deuchman gage) Nov. 8, 1881; minimum observed, 2.00 ft (Deuchman gage) Nov. 28, 1891.

EXTREMES FOR CURRENT YEAR.--Maximum daily mean gage height, 3.77 ft, June 19; minimum recorded, 1.62 ft, Feb. 29.

GAGE HEIGHT, FEET
WATER YEAR OCTOBER 2003 TO SEPTEMBER 2004
DAILY MEAN VALUES

DAY	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP
1	2.57	2.46	2.39	2.46	2.08	1.64	2.02	2.23	3.15	3.46	2.97	3.01
2	2.56	2.47	2.44	2.46	2.07	1.71	2.01	2.26	3.23	3.40	2.99	3.01
3	2.52	2.57	2.43	2.46	2.07	1.74	2.00	2.29	3.22	3.34	3.02	3.01
4	2.57	2.58	2.44	2.49	2.05	1.74	2.10	2.26	3.21	3.27	3.04	3.00
5	2.57	2.57	2.45	2.48	2.02	1.87	2.11	2.36	3.19	3.28	3.01	2.99
6	2.54	2.54	2.46	2.47	2.03	1.94	2.11	2.36	3.17	3.20	2.98	2.95
7	2.53	2.47	2.45	2.46	2.01	1.97	2.13	2.42	3.15	3.13	2.99	3.00
8	2.53	2.48	2.45	2.44	1.99	2.00	2.13	2.54	3.12	3.08	2.97	3.01
9	2.52	2.44	2.47	2.43	1.97	2.00	2.13	2.69	3.16	3.06	2.94	2.98
10	2.52	2.44	2.53	2.43	1.95	2.01	2.17	2.75	3.19	3.00	2.95	2.95
11	2.49	2.46	2.52	2.41	1.94	2.04	2.17	2.78	3.46	2.95	2.94	2.93
12	2.51	2.37	2.56	2.40	1.92	2.06	2.13	2.77	3.55	2.90	2.96	2.95
13	2.51	2.49	2.52	2.39	1.90	2.05	2.10	2.82	3.62	2.89	2.93	2.94
14	2.54	2.52	2.50	2.37	1.88	2.08	2.08	2.87	3.68	2.93	2.93	2.91
15	2.51	2.51	2.48	2.36	1.86	2.09	2.06	2.89	3.72	2.95	2.92	2.88
16	2.50	2.52	2.47	2.35	1.84	2.08	2.02	2.84	3.71	2.95	2.90	2.90
17	2.48	2.53	2.47	2.35	1.82	2.07	2.01	2.79	3.76	2.96	2.93	2.92
18	2.46	2.54	2.46	2.35	1.80	2.07	1.98	2.82	3.76	2.93	2.92	2.89
19	2.47	2.56	2.45	2.32	1.77	2.07	1.90	2.79	3.77	2.91	2.93	2.87
20	2.43	2.56	2.44	2.30	1.77	2.05	2.02	2.81	3.75	2.92	2.92	2.84
21	2.49	2.58	2.43	2.28	1.77	2.06	1.90	2.87	3.75	2.91	2.92	2.82
22	2.48	2.59	2.42	2.26	1.75	2.02	1.98	2.88	3.73	2.94	2.87	2.84
23	2.46	2.66	2.42	2.25	1.75	1.99	1.96	3.02	3.72	2.94	2.92	2.83
24	2.46	2.63	2.43	2.24	1.73	1.97	2.03	3.07	3.69	2.93	2.89	2.78
25	2.46	2.66	2.43	2.22	1.70	1.96	1.99	3.06	3.68	2.90	2.89	2.81
26	2.45	2.64	2.42	2.21	1.67	1.96	1.95	3.03	3.65	2.90	2.91	2.80
27	2.45	2.60	2.41	2.19	1.65	1.95	2.05	3.01	3.63	2.91	3.01	2.80
28	2.40	2.53	2.43	2.17	1.63	1.94	2.07	2.99	3.59	2.90	3.05	2.81
29	2.43	2.53	2.44	2.15	1.62	1.97	2.11	2.96	3.54	2.90	3.03	2.77
30	2.45	2.44	2.46	2.13	---	2.02	2.19	2.98	3.50	2.94	2.99	2.76
31	2.43	---	2.45	2.10	---	2.02	---	3.00	---	2.94	3.01	---
MEAN	2.49	2.53	2.46	2.33	1.86	1.97	2.05	2.75	3.50	3.02	2.96	2.90
MAX	2.57	2.66	2.56	2.49	2.08	2.09	2.19	3.07	3.77	3.46	3.05	3.01
MIN	2.40	2.37	2.39	2.10	1.62	1.64	1.90	2.23	3.12	2.89	2.87	2.76

04084255 LAKE WINNEBAGO NEAR STOCKBRIDGE, WI

LOCATION.--Lat 44°04'17", long 88°19'52", Stockbridge Indian Reservation, Calumet County, Hydrologic Unit 04030203, on east shore of Lake Winnebago, 300 ft south of County Highway E and 1.6 mi west of Stockbridge.

DRAINAGE AREA.--5,880 mi², at lake outlet at Menasha Dam. Area of Lake Winnebago, 215 mi².

PERIOD OF RECORD.--November 1982 to current year.

GAGE.--Water-stage recorder. Datum of gage is 745.05 ft above mean tide of New York City (levels by U. S. Army Corps of Engineers).

REMARKS.--Lake elevations controlled by dams at Menasha and Neenah, which are operated in the interest of navigation. Crests of both dams are at elevation 746.73 ft. Present limits of regulation are from 21 1/4 in. above the crest of Menasha dam to crest during navigation season, plus additional 18 in. below crest during winter. Data-collection platform and gage-height telemeter at station.

EXTREMES FOR PERIOD OF RECORD.--Maximum daily mean gage height, 3.85 ft, July 9, 11, 1993; minimum observed, 0.30 ft, Mar. 1, 1986.

EXTREMES FOR CURRENT YEAR.--Maximum daily mean gage height, 3.81 ft, June 20; minimum recorded, 1.57 ft, Feb. 29.

GAGE HEIGHT, FEET
WATER YEAR OCTOBER 2003 TO SEPTEMBER 2004
DAILY MEAN VALUES

DAY	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP
1	2.57	2.41	2.55	2.40	2.03	1.59	1.93	2.15	3.16	3.42	2.97	2.98
2	2.54	2.40	2.40	2.41	2.01	1.67	1.95	2.16	3.17	3.33	2.99	2.98
3	2.60	2.36	2.37	2.45	2.02	1.69	1.95	2.24	3.18	3.27	2.95	2.98
4	2.53	2.52	2.38	2.42	1.99	1.69	2.02	2.28	3.17	3.27	2.92	2.98
5	2.49	2.60	2.37	2.46	1.97	1.84	2.04	2.29	3.16	3.19	2.93	2.97
6	2.48	2.57	2.39	2.44	1.98	1.90	2.06	2.31	3.14	3.16	2.96	3.03
7	2.48	2.53	2.39	2.41	1.97	1.96	2.08	2.25	3.15	3.13	2.97	3.00
8	2.48	2.43	2.40	2.39	1.93	1.96	2.07	2.45	3.12	3.10	2.96	2.91
9	2.46	2.43	2.40	2.38	1.92	1.96	2.10	2.63	3.08	3.02	3.01	2.93
10	2.46	2.41	2.44	2.37	1.90	1.96	2.09	2.72	3.06	2.96	3.05	2.94
11	2.46	2.43	2.59	2.36	1.89	2.01	2.06	2.71	3.29	2.91	3.02	2.94
12	2.54	2.52	2.51	2.36	1.87	2.01	2.04	2.73	3.51	2.87	2.90	2.93
13	2.49	2.61	2.46	2.34	1.84	1.99	2.03	2.79	3.60	2.88	2.88	2.91
14	2.48	2.49	2.45	2.32	1.82	2.05	2.02	2.82	3.66	2.91	2.89	2.91
15	2.53	2.45	2.42	2.30	1.81	2.05	1.98	2.83	3.68	2.93	2.88	2.95
16	2.49	2.46	2.44	2.29	1.79	2.03	1.95	2.80	3.69	2.91	2.89	2.94
17	2.46	2.46	2.43	2.30	1.77	2.02	1.93	2.79	3.72	2.87	2.91	2.86
18	2.45	2.49	2.42	2.30	1.74	2.02	2.03	2.75	3.77	2.89	2.95	2.85
19	2.40	2.56	2.41	2.28	1.72	2.01	2.01	2.72	3.76	2.91	2.95	2.83
20	2.42	2.54	2.38	2.25	1.73	2.02	1.85	2.75	3.81	2.89	2.91	2.83
21	2.43	2.49	2.37	2.23	1.73	2.01	1.89	2.73	3.77	2.90	2.89	2.83
22	2.41	2.40	2.37	2.22	1.69	1.97	1.88	2.81	3.76	2.92	2.91	2.82
23	2.41	2.65	2.38	2.20	1.69	1.93	1.91	2.90	3.76	2.88	2.81	2.80
24	2.41	2.89	2.38	2.19	1.67	1.90	1.88	3.04	3.77	2.86	2.82	2.86
25	2.42	2.73	2.38	2.17	1.65	1.89	1.94	3.03	3.72	2.86	2.86	2.81
26	2.45	2.61	2.37	2.15	1.62	1.91	2.05	3.02	3.66	2.86	2.88	2.78
27	2.41	2.56	2.35	2.15	1.60	1.89	2.04	2.97	3.62	2.87	2.99	2.75
28	2.44	2.58	2.37	2.13	1.58	1.89	2.08	2.93	3.61	2.90	2.95	2.68
29	2.44	2.56	2.39	2.10	1.57	1.95	2.12	2.89	3.56	2.92	2.96	2.74
30	2.35	2.49	2.41	2.08	---	1.91	2.10	2.81	3.50	2.91	2.98	2.74
31	2.40	---	2.41	2.06	---	1.92	---	3.02	---	2.94	2.97	---
MEAN	2.46	2.52	2.41	2.29	1.81	1.92	2.00	2.69	3.49	2.99	2.93	2.88
MAX	2.60	2.89	2.59	2.46	2.03	2.05	2.12	3.04	3.81	3.42	3.05	3.03
MIN	2.35	2.36	2.35	2.06	1.57	1.59	1.85	2.15	3.06	2.86	2.81	2.68

WISCONSIN DISTRICT PUBLICATIONS PERTAINING TO LAKES

The reports published in a U.S. Geological Survey series are for sale by the U.S. Geological Survey, Box 25286, Federal Center, Denver, CO 80225. Prepayment is required. Remittance should be sent by check or money order payable to the U.S. Geological Survey. For more information and prices on USGS products, contact representatives at 1-888-ASK-USGS. For access to the new Publications Warehouse, go to <http://pubs.usgs.gov/> or link from the complete list of USGS publications and products at <http://www.usgs.gov/pub/prod/>. Reprints of journal articles may be obtained by writing directly to the author at U.S. Geological Survey, Wisconsin Water Science Center, 8505 Research Way, Middleton, WI 53562.

2004

Dupre, D.H., and Robertson, D.M., 2004, Water quality of Nippersink Creek and Wonder Lake, McHenry County, Illinois, 1994-2001: U.S. Geological Survey Scientific Investigations Report 2004-5085.

Rose, W.J., Robertson, D.M., and Mergener, E.A., 2004, Water Quality, Hydrology, and the Effects of Changes in Phosphorus Loading to Pike Lake, Washington County, Wisconsin, with Special Emphasis on Inlet-to-Outlet Short-Circuiting: U.S. Geological Survey Scientific Investigations Report 2004-5141, 32 p.

2003

Dunning, C.P., Thomas, J.C., and Lin, Y-F., 2003, Simulation of the Shallow Aquifer in the Vicinity of Silver Lake, Washington County, Wisconsin, Using Analytic Elements: U.S. Geological Survey Water-Resources Investigations Report 02-4204, 29 p.

Fitzpatrick, F.A., Garrison, P.J., Fitzgerald, S.A., and Elder, J.F., 2003, Nutrient, Trace-Element, and ecological history of Musky Bay, Lac Courte Oreilles, Wisconsin, as inferred from sediment cores: U.S. Geological Survey Water-Resources Investigations Report 02-4225, 141 p.

Fitzpatrick, F.A., and Peppler, M.C., 2003, Sedimentation and sediment chemistry, Neopit Mill Pond, Menominee Indian Reservation, Wisconsin, 2001: U.S. Geological Survey Open-File Report 03-23, 58 p.

Garn, H.S., Elder, J.F., and Robertson, D.M., 2003, Why study lakes? An overview of U.S. Geological Survey lake studies in Wisconsin: U.S. Geological Survey Fact Sheet FS-063-03, 8 p.

Graczyk, D.G., Hunt, R.J., Greb, S.R., Buchwald, C.A., and Krohelski, J.T., 2003, Hydrology, water quality, and yields, from near-shore flows to four lakes in northern Wisconsin, 1999-2001, U.S. Geological Survey Water-Resources Investigations Report 03-4144, 64 p.

Hunt, R.J., 2003, Ground water-lake interaction modeling using the LAK3 Package for MODFLOW2000: Ground Water, vol. 41, no. 2, p. 114-118.

- Hunt, R.J., Haitjema, H.M., Krohelski, J.T., and Feinstein, D.T., 2003, Simulating ground water-lake interactions: approaches and insights: *Ground Water*, vol. 41, no. 22, p. 227-237.
- Hunt, R.J., Pint, C.D., and Anderson, M.P., 2003, Using Diverse Data Types to Calibrate a Watershed Model of the Trout Lake Basin, Northern Wisconsin, p.600-604, In: MODFLOW and More 2003: Understanding through Modeling: Proceedings of the 5th International Conference of the International Ground Water Modeling Center. Golden, CO: Colorado School of Mines.
- Hunt, R.J., Saad, D.A., and Chapel, D.M., 2003, Numerical Simulation of Ground-Water Flow in La Crosse County, Wisconsin and into Nearby Pools of the Mississippi River. U.S. Geological Survey Water-Resources Investigations Report 03-4154. 36 p.
- John, R., Pint, C.D., Anderson, M.P., and Hunt, R.J., 2003, The Effects of Potential Climate Change on Lake Levels and Lake Capture Zones, p. 212-216, In MODFLOW and More 2003: Understanding through Modeling: Proceedings of the 5th International Conference of the International Ground Water Modeling Center. Golden, CO: Colorado School of Mines.
- Magnuson, J.J., Krohelski, J.T., Kunkel, K.E., and Robertson, D.M., 2003, Wisconsin's waters and climate: historical changes and possible futures: In Wisconsin's waters: a confluence of perspectives. *Transactions of the Wisconsin Academy*, p. 23-36.
- Pint, C.D., Hunt, R.J., and Anderson, M.P., 2003, Flow path delineation and ground water age, Allequash Basin, Wisconsin: *Ground Water*, vol. 41, no. 7, p. 895-902.
- Robertson, D.M., Rose, W.J., and Garn, H.S., 2003, Water quality and the effects of changes in phosphorus loading, Red Cedar Lakes, Barron and Washburn Counties, Wisconsin: U.S. Geological Survey Water-Resources Investigations Report 03-4238, 42 p.
- Robertson, D.M., Rose, W.J., and Saad, D.A., 2003, Water quality and the effects of changes in phosphorus loading to Muskellunge Lake, Vilas County, Wisconsin: U.S. Geological Survey Water-Resources Investigations Report 03-4011, 18 p.
- 2002
- Anderson, M.P., Hunt, R.J., Krohelski, J.T., and Chung, K., 2002, Using high hydraulic conductivity nodes to simulate seepage lakes: *Ground Water* 40(2), p. 119-124.
- Garn, H. S., 2002, Effects of lawn fertilizer on nutrient concentration in runoff from lakeshore lawns, Lauderdale Lakes, Wisconsin: USGS Water-Resources Investigations Report 02-4130, 6 p.
- Kelson, V.A., Hunt, R.J., and Haitjema, H.M., 2002, Improving a regional model using reduced complexity and parameter estimation: *Ground Water* 40(2), p. 138-149.
- Krohelski, J.T., Lin, Y., Rose, W.J., and Hunt, R.J., 2002, Simulation of Fish, Mud and Crystal Lakes and the shallow ground-water system, Dane County, Wisconsin: USGS Water-Resources Investigations Report 02-20.

Krohelski, J.T., Rose, W.J., and Hunt, R.J., 2002, Hydrologic investigation of Powell Marsh and its relationship to Dead Pike Lake Vilas County, Wisconsin: USGS Water-Resources Investigations Report 02-4034.

Madenjian, C.P., Fahnstiel, G.L., Johengen, T.H., Nalepa, T.F., Venderploeg, H.A., Fleischer, G.W., Schneeberger, P.J., Benjamin, D.M., Smith, E.B., Bence, J.R., Rutherford, E.S., Lavis, D.S., Robertson, D.M., Jude, D.J., and Ebener, M.P., 2002, Dynamics of the Lake Michigan food web, 1970-2000: Canadian Journal of Fisheries and Aquatic Sciences, Vol. 59, p. 736-753.

Robertson, D.M., Goddard, G.L., Mergerer, E.A., Rose, W.J., and Garrison, P.J., 2002, Hydrology and water quality of Geneva Lake, Walworth County, Wisconsin: U.S. Geological Survey Water Resources Investigations Report 02-4039, 73 p.

Robertson, D.M., and Lenz, B.N., 2002, Response of the St. Croix River Pools, Wisconsin and Minnesota, to various phosphorus-loading scenarios: U.S. Geological Survey Water-Resources Investigations Report 02-4181, 36 p.

2001

Hunt, R.J., Bradbury, K.R., and Krohelski, J.T., 2001, The effects of large-scale pumping and diversion on the water resources in Dane County, Wisconsin: USGS Fact Sheet FS-127-01, 4 p.

Hunt, R.J., Haitjema, H.M., Krohelski, J.T., and Feinstein, D.T., 2001, Simulating groundwater-lake interactions with models: MODFLOW and analytic element approaches. p. 328-334 in MODFLOW 2001 and Other Modeling Odysseys: Proceedings of the 4th International Conference of the International Ground Water Modeling Center. Golden, CO: Colorado School of Mines.

Lin, Y., Krohelski, J.T., and Hunt, R.J., 2001, Simulation of lake stage in two seepage lakes in southcentral Wisconsin using the LAK3 package for MODFLOW. p. 411-417 in MODFLOW 2001 and Other Modeling Odysseys: Proceedings of the 4th International Conference of the International Ground Water Modeling Center. Golden, CO: Colorado School of Mines.

2000

Elder, J.F., Robertson, D.M., and Garrison, P.J., 2000, Chemical composition of surficial deposits in Geneva Lake, Wisconsin: U.S. Geological Survey Fact Sheet FS-121-00.

Elder J.F., Rybicki, N.B., Carter, V.P., and Weintraub, V., 2000, Sources and yields of dissolved carbon in northern Wisconsin stream catchments with differing amounts of peatland: Wetlands, Vol. 20, No. 1, p.113-125.

Hunt, R.J., Graczyk, D.J., and Rose, W.J., 2000, Water flows in the Necedah National Wildlife Refuge: U.S. Geological Survey Fact Sheet FS-068-00, 4 p.

Hunt, R.J., Lin, J., Krohelski, J.T., and Juckem, P.F., 2000, Simulation of the shallow hydrologic system in the vicinity of Middle Genesee Lake, Wisconsin, using analytic

elements and parameter estimation: U.S. Geological Survey Water Resources Investigations Report 00-4136, 16 p.

Lathrop, R.C., Carpenter, S.R., and Robertson, D.M., 2000, Interacting factors causing exceptional water clarity in Lakes Mendota and Monona, Wisconsin, Proc. of the International Limnological Society, SIL, Dublin, Ireland, August, 1998.

Magnuson, J.J., Robertson, D.M. Wynne, R.H., Benson, B.J., Livigstone, D.M., Arai, T., Assel, R.A., Barry, R.G., Card, V., Kuusisto, E., Granin, N.G., Prowse, T.D., Stewart, K.M., and Vuglinski, V.S., 2000, Historical Trends in Lake and River Ice Cover in the Northern Hemisphere: Science, Vol. 289, No. 5485, p. 1743-1746.

Magnuson, J.J., Wynne, R.H., Benson, B.J., and Robertson, D.M., 2000, Lake and river ice as a powerful indicator of past and present climates, Proc. of the International Limnological Society, SIL, Dublin, Ireland, August, 1998.

Robertson, D.M., 2000, One-dimensional simulation of stratification and dissolved oxygen in McCook Reservoir, Illinois: U.S. Geological Survey Water Resources Investigations Report 00-4258, 17 p.

Robertson, D.M., Goddard, G.L., Helsel, D.R., and MacKinnon, K.L., 2000, Rehabilitation of Delavan Lake, Wisconsin: Lake and Reservoir Management, Vol. 20, No. 3, p. 155-176.

Robertson, D.M., and Rose, W.J., 2000, Hydrology, water quality, and phosphorus loading of Little St. Germain Lake, Vilas County, Wisconsin: U.S. Geological Survey Water Resources Investigations Report 00-4209, 8 p.

Robertson, D.M., Wynne, R.H., and Chang, W.Y.B., 2000, Variability in ice cover across the northern hemisphere during the 1900's associated with El Nino events, Proceedings of the International Limnological Society, SIL, Dublin, Ireland, August, 1998.

Saad, D.A., and Robertson, D.M., 2000, Water-resources-related information for the St. Croix Reservation and vicinity, Wisconsin: U.S. Geological Survey Water Resources Investigations Report 00-4133, 65 p.

Johnson, G.P., Hornewer, N.J., Robertson, D.M., and Olson, D.T., 2000, Methodology, data collection, and data analysis for determination of water-mixing patterns induced by aerators and mixers, U.S. Geological Survey Water Resources Investigations Report 00-4101, 72 p.

Grannemann, N.G., Hunt, R.J., Nicholas, J.R. Reilly, T.E., and Winter, T.C., 2000, The importance of ground water to the Great Lakes Region: U.S. Geological Survey Water Resources Investigations Report 00-4008, 12 p.

1999

Lathrop, R.C., Carpenter, S.R., and Robertson, D.M., 1999, Summer water clarity responses to phosphorus, Daphnia grazing, and internal mixing in Lake Mendota, Limnology and Oceanography, Vol. 44 (1), p. 137-146.

Krohelski, J.T., Feinstein, D.T., and Lenz, B.N., 1999, Simulation of stage and hydrologic budget for Shell Lake, Washburn County, Wisconsin, U.S. Geological Survey Water-Resources Investigations Report 99-4209, 23 p.

Panuska, J.C., and Robertson, D.M., 1999, Estimating phosphorus concentrations following alum treatment using apparent settling velocities, Lakes and Reservoir Management, Vol. 15 (1), p. 28-38.

1998

Hunt, R.J., Anderson, M.P., and Kelson, V.A., 1998, Improving a complex finite-difference ground water flow model through the use of an analytic element screening model, Ground Water, Vol. 36, No. 6, p. 1011-1017.

Hunt, R.J., Anderson, M.P., and Kelson, V.A., 1998, Linking an analytic element flow code to MODFLOW - Implementation and benefits. p. 477-504 in MODFLOW'98: Proceedings of the 3rd International Conference of the International Ground Water Center. Golden, CO: Colorado School of Mines.

Krabbenhoft, D.P., Gilmour, C.C., Benoit, J.M., Babiarz, C.L., Andren, A.W., and Hurley, J.P., 1998, Methyl mercury dynamics in littoral sediments of a temperate lake, Canadian Journal of Fisheries and Aquatic Sciences, Vol. 55, p. 835-844.

Robertson, D.M., Elder, J.F., Goddard, G.L., and James, W.F., 1998, Dynamics in phosphorus retention in wetlands upstream of Delavan Lake, Wisconsin, Lakes and Reservoir Management, Vol. 14, No. 4, p. 466-477.

Rose, W.J., and Robertson, D.M., 1998, Hydrology, water quality, and phosphorus loading of Kirby Lake, Barron County, Wisconsin, U.S. Geological Survey Fact Sheet FS-066-98, 4 p.

Walker, J.F. and Krabbenhoft, D.P., 1998, Groundwater and surface-water interactions in riparian and lake-dominated systems, in Kendall, C. and McDonnell, J.J. eds., Isotope tracers in catchment hydrology, Elsevier Publishing, New York, 839 P.

1997

Elder, J.F., Manion, B.J., and Goddard, G.L., 1997, Mesocosm experiments to assess factors affecting phosphorus retention and release in an extended Wisconsin wetland: U.S. Geological Survey Water-Resources Investigations Report 97-4272, 14 p.

Goddard, G.L., and Elder, J.F., 1997, Retention of sediments and nutrients in Jackson Creek Wetland near Delavan Lake, Wisconsin, 1993-95, U.S. Geological Survey Water-Resources Investigations Report 97-4014, 22 p.

Hornewer, N.J., Johnson, G.P., Robertson, D.M. and Hondzo, M., 1997, Field-scale tests for determining mixing patterns associated with coarse-bubble air diffuser configurations, Egan Quarry, Illinois, In: Environmental and Coastal Hydraulics: Protecting the Aquatic Habitat, Proceedings of the International Association of Hydraulic Research, San Francisco, CA, USA, p. 57-63.

Robertson, D.M., 1997, Regionalized loads of sediment and phosphorus to Lakes Michigan and Superior - High flow and long-term average, *Journal of Great Lakes Research*, Vol. 23, p. 416-439.

1996

Anderson, W.L., Robertson, D.M., and Magnuson, J.J., 1996, Evidence of recent warming and El Nino-related variation in ice breakup of Wisconsin lakes, *Limnology and Oceanography*, Vol. 41, p. 815-821.

Elder, J.F. and Goddard, G.L., 1996, Sediment and nutrient trapping efficiency of a constructed wetland near Delavan Lake, Wisconsin, 1993-1995: U.S. Geological Survey Fact Sheet FS-232-96.

Garn, H.S., Olson, D.L., Seidel, T.L., and Rose, W.J., 1996, Hydrology and water quality of Lauderdale Lakes, Walworth County, Wisconsin, 1993-94: U.S. Geological Survey Water-Resources Investigations Report 96-4235, 29 p.

Hunt, R.J. and Krohelski, J.T., 1996, The application of an analytical element model to investigate groundwater-lake interactions at Pretty Lake, Wisconsin, *Lake and Reservoir Management*, Vol. 12, p. 487-495.

Imberger, J., Robertson, D.M., and Boland, K., 1996, Lake Number: A quantitative indicator of mixing to be used in water quality management. *Scientific Impeller*, Solna, Sweden, No. 4, p. 9-15.

Kammerer, P.A., Jr., 1996, Hydrology and water quality of Park Lake, South-central Wisconsin: U.S. Geological Survey Fact Sheet FS-197-96.

Robertson, D.M., Field, S.J., Elder, J.F., Goddard, G.L., and James, W.F., 1996, Phosphorus dynamics of Delavan Lake Inlet in southeastern Wisconsin. U.S. Geological Survey Water-Resources Investigations Report 96-4160, 18 p.

1995

Assel, R.A. and Robertson, D.M. 1995, Changes in winter air temperatures near Lake Michigan during 1851 - 1993, as determined from regional lake-ice records, *Limnology and Oceanography*, v. 40, p 165-176.

Assel, R.A., Robertson, D.M., Hoff, M.H., and Selgeby, J.H., 1995, Climatic-change implications from long-term (1823-1994) ice records near the Laurentian Great Lakes: *Annals of Glaciology*, v. 21, p. 383-386.

Krabbenhoft, D.P., and Webster, K.E., 1995, Transient hydrogeological controls on the chemistry of a seepage lake: *Water Resources Research*, v. 31, no. 9, p. 2295-2305.

Krohelski, J.T. and Batten, W.G., 1995, Simulation of stage and the hydrologic budget of Devils Lake, Sauk County, Wisconsin: U.S. Geological Survey Open-File Report 94-348, 22 p.

Wentz, D.A., Rose, W.J., and Webster, K.E., 1995, Long-term hydrologic and biogeochemical responses of a soft water seepage lake in north central Wisconsin: Water Resources Research, v. 31, no. 1, p 199-212.

1994

Elder, J.F., 1994, Distribution and grain-size partitioning of metals in bottom sediments of an experimentally acidified lake: Water Resources Bulletin, v. 30, no. 2, p. 251-259.

Goddard, G.L., and Field, S.J., 1994, Hydrology and water quality of Whitewater and Rice Lakes in southeastern Wisconsin, 1990-91: U.S. Geological Survey Water-Resources Investigations Report 94-4101, 36 p.

Greb, S.R., and Wentz, D.A., 1994, Chemical budgets, in Klepinger, K.E., ed., RILWAS 1983-86: Wisconsin Regional Integrated Lake Watershed Acidification Study, Volume 1: Madison, Wisconsin Department of Natural Resources, PUBL-RS-909-94, Chapter 7, 20 p.

Hurley, J.P., Krabbenhoft, D.P., Babiarz, C.L., and Andren, A.W., 1994, Cycling processes of mercury across sediment/water interfaces in seepage lakes, in Baker, L.A. ed., Environmental Chemistry of Lakes and Reservoirs: Advances in Chemistry Series, American Chemical Society, Washington, D.C., p. 426-449.

Krabbenhoft, D.P., Bowser, C.J., Kendall, C., and Gat, J.R., 1994, Use of oxygen-18 and deuterium to assess the hydrology of ground-water/lake systems, in Baker, L.A. ed., Environmental Chemistry of Lakes and Reservoirs: Advances in Chemistry Series, American Chemical Society, Washington, D.C., p. 67-90.

Robertson, D.M., Anderson, W., and Magnuson, J.J., 1994, Relations between El Nino/Southern Oscillation events and the climate and ice cover of lakes in Wisconsin. P 48-57, in Greenland, D. ed., El Nino and Long-Term Ecological Research (LTER) Sites: Publication No. 18. LTER Network Office: University of Washington, Seattle, WA, USA., 57 p.

Robertson, D.M. and Imberger, J. 1994, Lake Number, a quantitative indicator of mixing used to estimate changes in dissolved oxygen, Internationale Revue der gesamten Hydrobiologie, v. 79, p. 159-176.

Watras, C.J., Bloom, N.S., Hurley, J.P., Fitzgerald, W.F., Andren, A.W., Krabbenhoft, D.P., and Porcella, D.B., 1994, Sources and fates of mercury and methylmercury in Wisconsin lakes, in Watras and Huckabee eds., Mercury as a Global Pollutant: Intergration and Synthesis, Lewis Pub., Chelsea, MI., p 153-177.

Wentz, D.A., 1994, Chemistry of snowpack and ground water, in Klepinger, K.E., ed., RILWAS 1983-86: Wisconsin Regional Integrated Lake Watershed Acidification Study, Volume 1: Madison, Wisconsin Department of Natural Resources, PUBL-RS-909-94, Chapter 6, 45 p.

Wentz, D.A., Krohelski, J.T., and Rose, W.J., 1994, Hydrology, in Klepinger, K.E., ed., RILWAS 1983-86: Wisconsin Regional Integrated Lake Watershed Acidification

Study, Volume 1: Madison, Wisconsin Department of Natural Resources, PUBL-RS-909-94, Chapter 7, 74 p.

1993

Field, S.J., 1993, Hydrology and water quality of Powers Lake, southeastern Wisconsin: U.S. Geological Survey Water-Resources Investigations Report 90-4126, 36 p.

Field, S.J., 1993, Hydrology and water quality of Wind Lake in southeastern Wisconsin: U.S. Geological Survey Water-Resources Investigations Report 91-4107, 61 p.

House, L.B., 1993, Simulation of the effects of hypothetical residential development on water levels in Graber Pond, Middleton, Wisconsin: U.S. Geological Survey Water-Resources Investigations Report 92-4029, 10 p.

House, L.B., Waschbusch, R.J., and Hughes, P.E., 1993, Water quality of an urban wet detention pond in Madison, Wisconsin, 1987-88: U.S. Geological Survey Open-File Report 93-172, 57 p.

Hughes, P.E., 1993, Hydrology, water quality, trophic status, and aquatic plants of Fowler Lake, Wisconsin: U.S. Geological Survey Water-Resources Investigations Report 91-4076, 44 p.

Rose, W.J., 1993, Hydrology of Little Rock Lake in Vilas County, north-central Wisconsin: U.S. Geological Survey Water-Resources Investigations Report 93-4139, 22 p.

Rose, W.J., 1993, Water and phosphorus budgets and trophic state, Balsam Lake, northwestern Wisconsin, 1987-1989: U.S. Geological Survey Water-Resources Investigations Report 91-4125, 28 p.

1992

Elder, J.F., Krabbenhoft, D.P., and Walker, J.F., 1992, Water, energy, and biogeochemical budgets (WEBB) program: Data availability and research at the northern temperate lakes site, Wisconsin: U.S. Geological Survey Open-File Report 92-48, 15 p.

Krabbenhoft, D.P., and Babiarz, C.L., 1992, Role of groundwater transport in aquatic mercury cycling: Water Resources Research, vol. 28, no. 12, p. 3119-3128.

Krabbenhoft, D.P., and Krohelski, J.T., 1992, Data on water quality, lake sediment, and lake-level fluctuation, St. Croix Indian Reservation, Wisconsin, 1981-87: U.S. Geological Survey Open-File Report 92-26, 53 p.

Robertson, D.M., Ragotzkie, R.A., and Magnuson, J.J., 1992, Lake ice records used to detect historical and future climatic changes: Climatic Change, v. 21, p. 407-427.

1991

Wentz, D.A., and Rose, W.J., 1991, Hydrology of Lakes Clara and Vandercook in North-Central Wisconsin: U.S. Geological Survey Water-Resources Investigations Report 89-4204, 24 p.

Watras, C.J., Andre, A.W., Bloom, N.S., Fitzgerald, W.F., Hurley, J.P., Krabbenhoft, D.P., Rada, R.G., Wiener, J.G., 1991, Mercury in temperate lakes: a mechanistic field study: *Verhandlungen Internat. Verein. Limnologie*, 24, p. 2199-2202.

Pre-1990

Walker, J.F., Pickard, S.A., and Sonzogni, W.C., 1989 Spreadsheet watershed modeling for nonpoint-source pollution management in a Wisconsin basin: *Water Resources Bulletin*, v. 25, no. 1, p. 139-147.

Wentz, D.A., Garrison, P.J., and Bockheim, J.G., 1989, Section 7—Chemical input-output budgets, in Knauer, D., and Brouwer, S.A., eds., *The Wisconsin Regional Integrated Lake-Watershed Acidification Study (RILWAS): 1981-1983*: Palo Alto, California, Electric Power Research Institute Report EA-6214, p. 7-1 to 7-30.

Wentz, D.A., and Rose, W.J., 1989, Interrelationships among hydrologic-budget components of a northern Wisconsin seepage lake and implications for acid-deposition modeling: *Archives of Environmental Contamination and Toxicology*, v. 18, p. 147-155.

Wentz, D.A., Rose, W.J., and Krohelski, J.T., 1989, Section 5—Hydrologic component, in Knauer, D., and Brouwer, S.A., eds., *The Wisconsin Regional Integrated Lake-Watershed Acidification Study (RILWAS): 1981-1983*: Palo Alto, California, Electric Power Research Institute Report EA-6214, p. 5-1 to 5-77.

Field, S.J., and Duerk, M.D., 1988, Hydrology and water quality of Delavan Lake in southeastern Wisconsin: U.S. Geological Survey Water-Resources Investigations Report 87-4168, 61 p.

Krug, W.R., Ostenson, N.A., and Krohelski, J.T., 1988, Prediction of the effects of mine dewatering on four lakes near Crandon, Wisconsin, by use of a water-budget model: U.S. Geological Survey Open-File Report 87-471, 63 p.

Wentz, D.A., Krohelski, J.T., Rose, W.J., Bockheim, J.G., Garrison, P.J., Knauer, D.R., and Goldstein, R.A., 1987, Hydrologic and chemical budgets of Wisconsin lakes, in Perrry, R., Harrison, R.M., Bel, J.N.B., and Lester, J.N., eds., *Acid Rain: Scientific and Technical Advances*, Selper Ltd., London, p. 309-316.

House, L.B., 1986, Stage fluctuations of Wisconsin Lakes: Wisconsin Geological and Natural History Survey Information Circular No. 49, 84 p.

House, L.B., 1984, Effects of urbanization on three ponds in Middleton, Wisconsin: U.S. Geological Survey Water-Resources Investigations Report 84-4051, 17 p.

Krug, W.R., and House, L.B., 1984, Evaluation of alternative reservoir-management practices in the Rock River basin, Wisconsin: U.S. Geological Survey Water-Resources Investigations Report 83-4186, 21 p.

House, L.B., 1981, An assessment of streamflow, water quality, and the effects of construction on impoundment on Bridge Creek at Augusta, Wisconsin: U.S. Geological Survey Water-Resources Investigations Open-File Report 81-1192, 25 p.

- Krug, W.R., 1981, Hydrologic effects of proposed changes in management practices, Winnebago Pool, Wisconsin: U.S. Geological Survey Water-Resources Investigations 80-107, 19 p.
- Batten, W.G., and Hindall, S.M., 1980, Sediment deposition in the White River Reservoir, northwestern Wisconsin: U.S. Geological Survey Water-Supply Paper 2069, 30 p.
- Novitzki, R.P., and Holmstrom, B.K., 1979, Monthly and annual water budgets of Lake Wingra, Madison, Wisconsin, 1971-77: U.S. Geological Survey Water-Resources Investigations 79-100, 31 p.
- Rose, W.J., 1977, Hydrologic considerations associated with dredging spring ponds in Wisconsin: U.S. Geological Survey Water-Resources Investigations 77-18, 35 p.
- Oakes, E.L., Hendrickson, G.E., and Zuehls, E.E., 1975, Hydrology of the Lake Wingra basin, Dane County, Wisconsin: U.S. Geological Survey Water-Resources Investigations 17-75, 31 p.
- Novitzki, R.P., 1971, Hydrologic investigations of Heart Lake, Green Lake County, Wisconsin: U.S. Geological Survey Administrative Report, 9 p.

APPENDIX

Wisconsin Lakes Team

Quality-Assurance Plan

Most lake studies and monitoring programs that are conducted by the USGS Wisconsin District entail water sampling and analysis to determine water quality and biological productivity. Because all sampling and analysis is subject to error and random variability, a certain proportion of the sampling effort should include quality-assurance samples. These samples are collected and/or prepared solely for the purpose of assessing the magnitude of error and random variability so that the accuracy and precision of all data can be evaluated. The plan for this quality-assurance sampling is described below.

Three types of QA/QC samples are collected:

blanks

provide information about accuracy and errors due to treatment or reagents

replicates

provide information about precision (variability)

standard additions (spikes)

provide information about accuracy and matrix interferences

Blank Sampling

B1. A **preservation blank** is prepared for each month of lake sampling. This consists of deionized water or organic blank water, to which is added any reagents or preservatives that are normally added to natural water samples. The blank is not taken to the field, but is shipped to the laboratory for analysis along with the natural water samples.

This blank sample is analyzed for the Nutrient Group¹ and chlorophyll-a.

B2. At one randomly-chosen lake each month, a **field blank** is prepared. This consists of deionized water or inorganic blank water treated exactly the same as regular samples. During winter, the field blank is analyzed for total phosphorus (TP) only; during summer, it is analyzed for TP and chlorophyll-a, and in the spring, it is analyzed for the Nutrient Group and chlorophyll-a.

¹ Nutrient Group = all phosphorus and nitrogen species that are commonly determined in lakes (total phosphorus, nitrate + nitrite, ammonia, total Kjeldahl nitrogen, total nitrogen)

Replicate Sampling

R1. At all lakes in program, **triplicate samples** are taken near water surface in summer for analysis of total phosphorus and chlorophyll-a. At two of these lakes, a set of triplicate samples is also taken from near-bottom water, for analysis of total phosphorus.

R2. At three selected lakes in the spring (different lakes each year), **triplicate samples** are taken near water surface for analysis of Nutrient Group.

R3. At one lake each year, **5 replicate samples** are taken near water surface for analysis of total phosphorus and chlorophyll-a.

Standard Addition Testing

S1. At Delavan Lake and one other lake (to be determined each year), **5 replicate samples** are taken in August for **a standard addition (spike) test**. The spike consists of addition of a prepared phosphorus solution (standard) of known volume and concentration, such that the expected result of analysis is the natural water TP concentration plus the known addition. One sample from each set will receive a low-concentration spike, one will receive a high-concentration spike, and three will receive no spike (the mean of these gives the natural water TP concentration).

Data and results of replicate sampling and field blank testing in water year 2000 are shown in Table A1.

Table A1. Analyses of replicate samples from Wisconsin lakes in water years 2001-2004. See text for procedures used. All data except chlorophyll in milligrams per liter. Symbol "<" indicates less than given detection limit (DL); mean and standard deviation not calculated for datasets containing values less than DL. All samples collected near surface (approximately 0.5 m depth) unless otherwise indicated.

Parameter	Lake	Date	Replicate Data			Percent		
						Standard Mean	Standard Deviation	
Total Phosphorus	Buffalo	7/23/01	0.276	0.275	0.277	0.276	0.001	0.4
	Delavan	7/15/01	0.027	0.027	0.031	0.028	0.002	8.2
	Delavan	8/19/01	0.031	0.027	0.035	0.031	0.004	12.9
	Geneva	7/15/01	0.005	<0.005	<0.005			
	Little Green	7/23/01	0.069	0.074	0.072	0.072	0.003	3.5
	Middle	6/17/01	0.012	0.012	0.017	0.016	0.014	0.003
	Muskego	4/18/01	0.039	0.044	0.047		0.043	0.004
	Muskego	7/25/01	0.030	0.031	0.031		0.031	0.001
	Oconomowoc	7/17/01	0.010	0.011	0.010		0.010	0.001
	Oconomowoc	8/23/01	0.011	0.010	0.009		0.010	0.001
	Okauchee	8/20/01	0.013	0.015	0.015		0.014	0.001
	Red Cedar	7/9/01	0.021	0.022			0.022	0.001
Total Phosphorus, near bottom	Delavan	7/15/02	0.026	0.026	0.027	0.031	0.028	0.002
	Geneva	7/16/02	0.008	0.008	0.008		0.008	0.000
	Little Muskego	7/1/02	0.016	0.016	0.017		0.016	0.001
	Potter	8/5/02	0.041	0.036	0.042	0.043	0.041	0.003
	Little St. Germain	7/22/02	0.061	0.060	0.059		0.060	0.001
	Delavan	4/14/03	0.057	0.057	0.057		0.057	0.000
	Delavan	8/12/03	0.044	0.043	0.041		0.043	0.002
	Lac La Belle	8/19/03	0.015	0.012	0.012		0.013	0.002
	Butternut	8/13/03	0.040	0.042			0.041	0.001
	Delavan	7/20/04	0.031	0.020	0.041		0.031	0.011
	Big Cedar	8/18/04	0.012	0.011	0.012		0.012	0.001
Dissolved Phosphorus	Geneva	7/15/01	0.017	0.020	0.021		0.019	0.002
	Red Cedar	7/9/01	0.187	0.228	0.262		0.226	0.038
	Wind	7/8/02	0.084	0.089	0.092		0.088	0.004
	Wind	8/19/03	0.194	0.192	0.165		0.184	0.016
	Delavan	7/15/01	0.010	<0.002	<0.007			
Dissolved Ammonia	Geneva	4/17/01	<0.002	<0.002				
	Oconomowoc	8/23/01	0.002	<0.002	<0.002			
	Delavan	4/14/03	0.022	0.023	0.023		0.023	0.001
	Delavan	7/15/01	0.026	0.013	0.021		0.020	0.007
	Geneva	4/17/01	0.014	0.022			0.018	0.006
Dissolved Ammonia	Muskego	4/18/01	0.086	0.083	0.084		0.084	0.002
	Oconomowoc	8/23/01	0.027	0.028	0.022		0.026	0.003
	Delavan	4/14/03	<0.015	<0.015	<0.015			

* Algal bloom on lake.

	Delavan	7/15/01	0.560	0.580	0.560	0.567	0.012	2.0
Total Kjeldahl Nitrogen	Geneva	4/17/01	0.390	0.390		0.390	0.000	0.0
	Muskego	4/18/01	1.200	1.100	1.200	1.167	0.058	4.9
	Oconomowoc	8/23/01	0.490	0.500	0.520	0.503	0.015	3.0
	Delavan	4/14/03	0.640	0.640	0.620	0.633	0.012	1.8

Parameter	Lake	Date	Replicate Data			Mean	Standard Deviation	Percent Standard Deviation
Dissolved Nitrate plus Nitrite	Delavan	7/15/01	0.014	0.008	0.007	0.010	0.004	39.2
	Geneva	4/17/01	0.113	0.115		0.114	0.001	1.2
	Muskego	4/18/01	0.102	0.103	0.104	0.103	0.001	1.0
	Oconomowoc	8/23/01	0.370	0.371	0.369	0.370	0.001	0.3
	Delavan	4/13/03	<0.022	<0.022	<0.022			

Chlorophyll-a (micrograms per liter)	Buffalo	7/23/01	14.0	16.0	17.0	15.7	1.5	9.8		
	Delavan	7/15/01	4.9	4.0	4.8	4.6	0.5	10.8		
	Geneva	7/15/01	<1.0	<1.0	1.1					
	Little Green	7/23/01	23.0	24.0	24.0	23.7	0.6	2.4		
	Middle	6/17/01	1.6	4.7		3.2	2.2	69.6		
	Muskego	7/25/01	6.6	3.2	3.2	4.3	2.0	45.3		
	Oconomowoc	7/17/01	2.6	2.8	2.3	2.6	0.3	9.8		
	Okawhee	8/20/01	8.0	8.0	8.0	8.0	0.0	0.0		
	Powers	7/25/01	4.8	5.0	5.5	5.1	0.4	7.1		
	Red Cedar	7/9/01	5.2	3.7		4.5	1.1	23.8		
	Delavan	7/15/02	9.7	6.9	8.0	8.1	8.2	1.2	14.1	
	Geneva	7/16/02	0.74	1.00	0.96		0.9	0.1	15.6	
	Little Muskego	7/1/02	1.74	1.50	1.34		1.5	0.2	13.2	
	Potter	8/5/02	10.8	10.3	11.9	9.77	11.0	10.8	0.8	7.4
	Little St. Germain	7/22/02	63.8	62.2	69.7		65.2	4.0	6.1	
	Lac La Belle	8/19/03	3.3	3.7	3.5		3.5	0.2	5.3	
	Butternut	8/13/03	44.00	46.10	45.20		45.1	1.1	2.3	
	Delavan	7/20/04	10.4	11.6	10.5		10.8	0.7	6.1	
	Big Cedar	8/18/04	8.36	8.56	8.61		8.51	0.13	1.6	

Table A2. Data from standard addition tests using stock solution containing 5.00 mg/L phosphorus. See text for detail of procedures. All concentration data in milligrams per liter.

Lake, Date	Original Sample Concentration	Stock Solution Volume Added (milliliters)	Final Expected Concentration	Actual Detected Concentration	Percent Recovery
Delavan 08/12/03	0.043	0.310	0.056	0.058	116%
	0.043	1.250	0.094	0.099	108%

Table A3. Data from tests of blanks, 2001-2004. All data in milligrams per liter, unless otherwise indicated. < = less than given detection limit; E = estimated value.

Delavan Lake. Analyses at USGS National Water Quality Laboratory, Lakewood, CO.

Parameter	2/19/01	4/17/01	7/15/01	2/21/02	4/17/02	7/14/02	2/20/03	4/16/03	6/22/04
Total P	E 0.003	< 0.004	< 0.004	< 0.004	< 0.004	< 0.004	0.006	E 0.002	< 0.004
Dissolved orthophosphate			< 0.007			< 0.007	< 0.007		< 0.007
Chlorophyll a	< 0.1	< 0.1		< 0.1		< 0.1	< 0.1		< 0.260
Chlorophyll b	< 0.1			< 0.1		< 0.1	< 0.1		< 0.260
Total Kjeldahl Nitrogen (as N)			< 0.08		< 0.10	E 0.05	E 0.05		< 0.10
Ammonia (as N)			< 0.02			0.037	0.034		< 0.015
Nitrate + Nitrite (as N)			< 0.05			E 0.011	0.008		< 0.022

Big Cedar Lake, south site, near West Bend, WI. Analyses at Wisconsin State Laboratory of Hygiene, Madison, WI

Parameter	4/22/02	8/8/02
Total P	< 0.005	< 0.005
Dissolved orthophosphate	< 0.002	
Chlorophyll a	< 1.00	< 0.26
Total Kjeldahl Nitrogen (as N)	< 0.14	
Ammonia (as N)	< 0.013	
Nitrate + Nitrite (as N)	< 0.010	
Calcium, dissolved	< 0.20	
Magnesium, dissolved	< 0.20	
Potassium, dissolved	< 1.0	
Sodium, dissolved	< 0.10	
Iron, dissolved (micrograms per liter)	< 100	
Manganese, dissolved (micrograms per liter)	< 1	

Little Cedar Lake, south site, near West Bend, WI. Analyses at Wisconsin State Laboratory of Hygiene, Madison, WI

Parameter	4/29/03
Total P	< 0.005
Dissolved orthophosphate	< 0.002
Total Kjeldahl Nitrogen (as N)	< 0.14
Ammonia (as N)	< 0.013
Nitrate + Nitrite (as N)	0.016

Little Cedar Lake, North site, near West Bend, WI. Analyses at Wisconsin State Laboratory of Hygiene, Madison, WI

Parameter	2/21/03
Total P	< 0.005